



Europäisches Patentamt
European Patent Office
Office européen des brevets



(11) **EP 1 344 820 A1**

(12) **EUROPEAN PATENT APPLICATION**
published in accordance with Art. 158(3) EPC

(43) Date of publication:
17.09.2003 Bulletin 2003/38

(51) Int Cl.7: **C12N 9/42**, C11D 3/386,
D06M 16/00, C12R 1/645,
C12N 1/15, D06M 101/06

(21) Application number: **01997189.4**

(22) Date of filing: **21.11.2001**

(86) International application number:
PCT/JP01/10188

(87) International publication number:
WO 02/042474 (30.05.2002 Gazette 2002/22)

(84) Designated Contracting States:
**AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU
MC NL PT SE TR**
Designated Extension States:
AL LT LV MK RO SI

- **BABA, Yuko, c/o Meiji Seika Kaisha, Ltd.**
Sakado-shi, Saitama 350-0289 (JP)
- **KOGA, Jinichiro, c/o Meiji Seika Kaisha, Ltd.**
Sakado-shi, Saitama 350-0289 (JP)
- **KUBOTA, Hidetoshi, c/o Meiji Seika Kaisha, Ltd.**
Sakado-shi, Saitama 350-0289 (JP)

(30) Priority: **21.11.2000 JP 2000354296**

(71) Applicant: **Meiji Seika Kaisha, Ltd.**
Tokyo 104-8002 (JP)

(74) Representative: **Gillard, Richard Edward**
Elkington and Fife
Prospect House
8 Pembroke Road
Sevenoaks
Kent TN13 1XR (GB)

(72) Inventors:
• **NAKANE, Akitaka, c/o Meiji Seika Kaisha, Ltd.**
Sakado-shi, Saitama 350-0289 (JP)

(54) **ZYGOMYCETES-ORIGIN ENDOGLUCANASE LACKING CELLULOSE-BINDING DOMAIN**

(57) This invention relates to a protein that is a *Zygomycetes*-derived endoglucanase lacking the cellulose-binding domain and exhibits endoglucanase activity, and a method for using the same. This invention can enhance effects of an endoglucanase enzyme on fabric treatment such as reduction of fuzzing, improvement in

feel and appearance, color clarification, partial color change, and softening of cellulose-containing fabrics and on performance improvement in the deinking of waste paper and drainage of paper pulp.

EP 1 344 820 A1

DescriptionTechnical Field

- 5 **[0001]** The present invention relates to an endoglucanase enzyme, the cellulose-binding domain of which has been deleted, with enhanced effects in the treatment of cellulose-containing fabrics and with applications regarding detergents or paper pulp, a method for producing the same, and a cellulase preparation with enhanced effects.

Background Art

- 10 **[0002]** Treatment of cellulose-containing fabrics with cellulase is carried out to provide the fabrics with desired properties. For example, treatment with cellulase is carried out in the fabric industry in order to improve the feel and appearance of cellulose-containing fabrics or to give colored cellulose-containing fabrics an appearance of "stone-washed" material, i.e., partial color change (European Patent No. 307,564).
- 15 **[0003]** Colored cellulose-containing fabrics are known to become fuzzy after repeated washings and to lose their vividness. Incorporation of cellulase into a detergent can remove fuzz and make the color of fabrics vivid, i.e., clarify the color (European Patent No. 220,016). Thus, detergents containing cellulase are commercially available mainly in Europe and America.
- 20 **[0004]** In the aforementioned application, cellulases derived from *Trichoderma* or *Humicola* (both are wood-rotting fungi) are mainly used. Recently used cellulase preparations are produced by isolating endoglucanases, which are highly active in fabric treatment, from these cellulase components and enhancing their effects with genetic engineering in order to improve commercial efficiency. Examples of these highly active endoglucanases include: *Humicola insolens*-derived EG V (WO 91/17243) and NCE4 (WO 98/03640) that strongly act on cotton fabrics; and *Rhizopus oryzae*-derived RCE I, RCE II, and RCE III, *Mucor circinelloides*-derived MCE I and MCE II, and *Phycomyces nitens*-derived
- 25 PCE I (WO 00/24879) that strongly act on lyocell fabrics.
- [0005]** Among endoglucanases used in the aforementioned applications, EG V (WO 91/17243), NCE4 (WO 98/03640), and RCE I, RCE II, RCE III, MCE I, MCE II, and PCE I (WO 00/24879) are presumed to belong to the same family (family 45) because of their amino acid sequences, and these enzymes have common structural properties. Specifically, each of these endoglucanases comprises a cellulose-binding domain for binding to cellulose as its substrate (hereinafter referred to as "CBD"), a catalytic active domain as an active center (hereinafter referred to as "CAD"),
- 30 and a linker domain with a high hydrophilic amino acid residue content for linking these two domains.
- [0006]** EG V, an endoglucanase belonging to family 45, was studied using an enzyme, the CBD domain of which had been deleted (JP Patent Publication (PCT Translation) No. 9-500667, Enzyme and Microbial Technology, 27 (2000), 325-329). However, no improvement has been reported regarding the activity of endoglucanase for removing fuzz from
- 35 cellulose fabrics through the deletion of the cellulose-binding domain (CBD). There are still many unclarified matters concerning the role of the cellulose-binding domain of endoglucanase in the exhibition of endoglucanase activity, and research thereof is limited to cellulase derived from a specific fungus, namely, *Trichoderma* (Kioyula, A. et al., *Trichoderma Gliocladium*, 2, (1998), 3-23). There has been no detailed research on the cellulose-binding domain of the *Zygomycetes*-derived endoglucanase.
- 40 **[0007]** Up to the present, several contrivances have been made in order to improve the effect or performance of cellulase in the above applications. For example, mutation was applied to an enzyme for improvement thereof, or culture conditions were modified in order to improve the productivity of the enzyme. Due to the high cost of cellulase to be used, however, the effect of cellulase should be further improved in order to provide a cellulase preparation that is worth using at an industrially practical level. Recently used cellulase preparations are produced by reinforcing only
- 45 endoglucanase, which is highly active in fabric treatment, with genetic engineering in order to improve commercial efficiency. Accordingly, it is desirable to further improve the activity of such highly active cellulase.

Disclosure of the Invention

- 50 **[0008]** An object of the present invention is to provide an endoglucanase with improved activity, a cellulase preparation comprising the same, and various methods for treating cellulose-containing fabrics using the same.
- [0009]** The present inventors have conducted concentrated studies on the *Zygomycetes*-derived endoglucanase concerning the role of a cellulose-binding domain when acting on cellulose-containing fabrics. As a result, they have found that an endoglucanase lacking the cellulose-binding domain has much higher activity of removing fuzz from
- 55 cotton, lyocell, or the like compared to an endoglucanase having a cellulose-binding domain. This led to the completion of the present invention.
- [0010]** Specifically, the present invention relates to a *Zygomycetes*-derived endoglucanase that has enhanced effects for removing fuzz from cellulose-containing fabrics (e.g., an enzyme comprising amino acid sequences of RCE I, RCE

II, RCE III, MCE I, MCE II, and PCE I, which attained enhanced effects of removing fuzz from cellulose-containing fabrics through the deletion of the cellulose-binding domain, and exhibiting endoglucanase activity, a modified protein thereof exhibiting endoglucanase activity, or a homologue of the protein or the modified protein) and a cellulase preparation comprising such endoglucanase. The present invention also relates to an endoglucanase that was produced in a host cell transformed with a gene encoding such an endoglucanase, and further relates to a method for treating cellulose-containing fabrics using the endoglucanase, which attained improved activity through the deletion of the cellulose-binding domain, or the cellulase preparation.

[0011] More specifically, the present invention includes the following.

- (1) A protein that is a *Zygomycetes*-derived endoglucanase lacking the cellulose-binding domain and exhibits endoglucanase activity.
- (2) A protein that is a *Zygomycetes*-derived endoglucanase belonging to family 45 lacking the cellulose-binding domain and exhibits endoglucanase activity.
- (3) The protein according to (1) or (2), wherein the *Zygomycetes* are microorganisms selected from the group consisting of those belonging to *Rhizopus*, *Mucor*, and *Phycomyces*.
- (4) The protein according to (3), wherein the *Zygomycetes* are microorganisms belonging to *Rhizopus*.
- (5) A protein comprising an amino acid sequence as shown in SEQ ID NO: 1, 3, 5, 7, 9, or 11, wherein the cellulose-binding domain has been deleted therefrom, and exhibiting endoglucanase activity, a modified protein thereof exhibiting endoglucanase activity, or a homologue of the protein or the modified protein exhibiting endoglucanase activity.
- (6) A protein comprising an amino acid sequence as shown in SEQ ID NO: 1, 3, 5, 7, 9, or 11, wherein the cellulose-binding domain has been deleted therefrom, and exhibiting endoglucanase activity.
- (7) A gene encoding the protein, modified protein thereof, or homologue of the protein or the modified protein according to any one of (1) to (6).
- (8) An expression vector comprising the gene according to (7).
- (9) A host cell transformed with the expression vector according to (8).
- (10) The host cell according to (9), which is a filamentous fungus.
- (11) The host cell according to (10), which is a microorganism belonging to *Humicola*.
- (12) A method for producing a protein comprising steps of culturing the host cell according to any one of (9) to (11) and collecting from the host cell obtained by the step of culturing or its culture product the protein, modified protein thereof, or homologue of the protein or the modified protein according to any one of (1) to (6).
- (13) A protein produced by the method according to (12).
- (14) A cellulase preparation comprising the protein, modified protein thereof, or homologue of the protein or the modified protein according to any one of (1) to (6) and (13).
- (15) A method for treating cellulose-containing fabrics comprising a step of bringing cellulose-containing fabrics into contact with the protein, modified protein thereof, or homologue of the protein or the modified protein according to any one of (1) to (6) and (13) or the cellulase preparation according to (14).
- (16) A method for reducing the rate at which cellulose-containing fabrics become fuzzy or reducing fuzzing in cellulose-containing fabrics comprising a step of bringing cellulose-containing fabrics into contact with the protein, modified protein thereof, or homologue of the protein or the modified protein according to any one of (1) to (6) and (13) or the cellulase preparation according to (14).
- (17) A method of weight loss treatment of cellulose-containing fabrics to improve the feel and appearance thereof comprising a step of bringing cellulose-containing fabrics into contact with the protein, modified protein thereof, or homologue of the protein or the modified protein according to any one of (1) to (6) and (13) or the cellulase preparation according to (14).
- (18) A method of color clarification of colored cellulose-containing fabrics comprising a step of treating colored cellulose-containing fabrics with the protein, modified protein thereof, or homologue of the protein or the modified protein according to any one of (1) to (6) and (13) or the cellulase preparation according to (14).
- (19) A method of providing colored cellulose-containing fabrics with partial color change comprising a step of treating colored cellulose-containing fabrics with the protein, modified protein thereof, or homologue of the protein or the modified protein according to any one of (1) to (6) and (13) or the cellulase preparation according to (14).
- (20) A method for reducing the rate at which cellulose-containing fabrics become stiff or reducing stiffness in cellulose-containing fabrics comprising a step of treating cellulose-containing fabrics with the protein, modified protein thereof, or homologue of the protein or the modified protein according to any one of (1) to (6) and (13) or the cellulase preparation according to (14).
- (21) The method according to any one of (15) to (20), wherein fabrics are treated through soaking, washing, or rinsing thereof.
- (22) An additive to detergent comprising the protein, modified protein thereof, or homologue of the protein or the

modified protein according to any one of (1) to (6) and (13) or the cellulase preparation according to (14) in a non-dusting granular form or a stabilized liquid form.

(23) A detergent composition comprising the protein, modified protein thereof, or homologue of the protein or the modified protein according to any one of (1) to (6) and (13) or the cellulase preparation according to (14).

5 (24) A method of deinking waste paper using a deinking agent wherein the protein, modified protein thereof, or homologue of the protein or the modified protein according to any one of (1) to (6) and (13) or the cellulase preparation according to (14) is used in a step of deinking waste paper.

(25) A method for improving drainage of paper pulp comprising a step of treating paper pulp with the protein, modified protein thereof, or homologue of the protein or the modified protein according to any one of (1) to (6) and
10 (13) or the cellulase preparation according to (14).

(26) A method for improving digestibility of animal feeds comprising a step of treating animal feeds with the protein, modified protein thereof, or homologue of the protein or the modified protein according to any one of (1) to (6) and (13) or the cellulase preparation according to (14).

15 1. Endoglucanase lacking cellulose-binding domain

[0012] The present invention relates to a protein that is a *Zygomycetes*-derived endoglucanase lacking the cellulose-binding domain and exhibits endoglucanase activity.

[0013] In this description, "*Zygomycetes*" refer to microorganisms belonging to *Zygomycota*, i.e., fungi that generate zygospores through gametangial copulation upon gametogony. The *Zygomycota* includes *Zygomycetes* and *Trichomycetes*. The *Zygomycetes* used in the present invention are not particularly limited. Microorganisms belonging to the *Zygomycetes* are preferable, those belonging to the *Mucorales* are more preferable, those belonging to *Rhizopus*,
20 *Mucor*, or *Phycomyces* are still more preferable, and those belonging to the *Rhizopus* are the most preferable.

[0014] In this description, "endoglucanase activity" refers to CMCase activity. Further, "CMCase activity" refers to an activity for hydrolyzing carboxymethylcellulose (CMC, Tokyo Kasei Kogyo, Japan), and one unit is defined as the amount of an enzyme which produces reducing sugars corresponding to 1 μ mol of glucose per minute by measuring
25 amounts of the reducing sugars released after incubation of a test protein with a CMC solution for a given period of time.

[0015] Endoglucanase activity can be determined by, for example, a procedure as described below. At the outset, 0.5 ml of a solution containing a test protein is added to 0.5 ml of 50 mM acetic acid-sodium acetate buffer solution
30 (pH 6.0) with 2% CMC dissolved therein, and the mixture is subjected to incubation at 50°C for 30 minutes. Subsequently, the concentration of the reducing sugars produced in the resulting reaction solution is quantified by the 3,5-dinitrosalicylic acid (DNS) method. Specifically, 3.0 ml of a DNS reagent is added to 1.0 ml of the reaction solution 30 minutes after the reaction, and the mixture is subjected to incubation in a boiling water bath for 5 minutes. Thereafter, the incubation product is diluted with 8.0 ml of distilled water, and the absorbance at 540 nm is measured. A calibration
35 curve is prepared using a gradually diluted glucose solution, and the amount of the reducing sugars produced in the enzyme reaction solution is determined. Activity is determined using an amount of the enzyme that produces reducing sugars corresponding to 1 μ mol glucose per minute as one unit. This DNS reagent can be prepared in accordance with publication such as Seibutsu Kagaku Jikkenhou 1 - Kagentou no Teiryohou (Biochemical Experimentation 1-
40 The method for quantifying reducing sugar) (p. 19-20, Sakuzo Fukui, Center for Academic Publications Japan), and can be prepared in the manner described below. At the outset, 880 ml of 1% 3,5-dinitrosalicylic acid solution and 255 g of Rochelle salt are added to 300 ml of an aqueous solution of 4.5% sodium hydroxide (solution A). Separately, 10 g of crystalline phenol is added to 22 ml of an aqueous solution of 10% sodium hydroxide, and water is further added and dissolved in the mixture to bring the amount thereof to 100 ml (solution B). Sodium bicarbonate (6.9 g) is added to 69 ml of solution B and dissolved therein, solution A is poured therein, and the mixture is stirred and mixed until the
45 Rochelle salt is thoroughly dissolved. The mixture is allowed to stand for 2 days and then filtered.

[0016] The term "endoglucanase" used herein refers to an enzyme exhibiting endoglucanase activity, i.e., endo-1,4- β -glucanase (EC 3. 2. 1. 4). This enzyme hydrolyzes the β -1,4-glucopyranosyl bond of β -1,4-glucan.

[0017] Endoglucanases are classified into several families based on information such as their amino acid sequences. The endoglucanase according to the present invention may belong to any family, and it preferably belongs to family
50 45. The endoglucanase "belonging to family 45" refers to those types having a consensus sequence, (Ser, Thr, or Ala)-Thr-Arg-Tyr-(Trp, Tyr, or Phe)-Asp-Xaa-Xaa-Xaa-Xaa-Xaa-(Cys or Ala), in the catalytic active domain (CAD). *Humicola insolens*-derived EG V (JP Patent Publication (PCT Translation) No. 5-509223), NCE4 (WO 98/03640), and the like also belong to family 45.

[0018] A protein of endoglucanase belonging to family 45 is comprised a catalytic active domain (CAD), a cellulose-binding domain (CBD), and a linker domain for linking based on function. The cellulose-binding domain (CBD) is known to exist as a domain linking to cellulose as its name suggests, and the conservation of the following consensus sequence is confirmed as a feature of the sequence (Hoffren, A. -M. et al., Protein Engineering 8: 443-450, 1995).

CBD consensus sequence:

1
 Xaa Xaa Xaa Xaa Xaa Xaa Gln Cys Gly Gly Xaa Xaa Xaa Xaa
 20
 Gly Xaa Xaa Xaa Cys Xaa Xaa Xaa Xaa Xaa Cys Xaa Xaa
 30
 Xaa Xaa Xaa Asn Xaa Xaa Tyr Xaa Gln Cys Xaa (SEQ ID NO: 17)

[0019] In this sequence, Xaa is independently any amino acid; and Xaas at positions 20, 21, 22, 23, 24, 30 and 31 may be independently absent. Other Xaas are always present and are independently any amino acids. Amino acids other than Xaa are expressed in three-letter abbreviations. CBD is linked to either the N-terminal side or C terminal side of CAD through a linker domain. Also reported is *Humicola insolens*-derived NCE5 (amino acid sequence: SEQ ID NO: 38, cDNA sequence: SEQ ID NO. 39) such as a family 45 endoglucanase that does not originally have CBD.

[0020] Although there is no definite recognition sequence for the linker domain, the sequence is rich in hydrophilic amino acid residues such as Ser or Thr, and its length varies depending on types of endoglucanases.

[0021] Examples of the *Zygomycetes*-derived endoglucanase according to the present invention include enzymes exhibiting endoglucanase activity derived from *Rhizopus*, *Phycomyces*, or *Mucor* described in WO 00/24879, i.e., RCE I (SEQ ID NO: 1), RCE II (SEQ ID NO: 3), RCE III (SEQ ID NO: 5), MCE I (SEQ ID NO: 7), MCE II (SEQ ID NO: 9), and PCE I (SEQ ID NO: 11). Locations of each of the domains in the amino acid sequence of these enzymes are as shown in Table 1 below.

Table 1

	CBD	A portion in linker domain	CAD
SEQ ID NO: 1	3 to 38	99 to 108	109 to 315
SEQ ID NO: 3	3 to 38	127 to 136	137 to 343
	50 to 85		
SEQ ID NO: 5	3 to 40	122 to 131	132 to 337
SEQ ID NO: 7	3 to 40	104 to 113	114 to 316
SEQ ID NO: 9	3 to 40	153 to 162	163 to 365
	52 to 89		
SEQ ID NO: 11	3 to 40	115 to 124	125 to 327

[0022] Amino acid sequences at the N-terminuses of RCE I, MCE I, and PCE I are respectively identified as shown in SEQ ID Nos: 14, 15, and 16 (WO 00/24879).

[0023] The protein according to the present invention should not comprise a cellulose-binding domain in the aforementioned endoglucanase. As long as the protein has endoglucanase activity, there is no particular limitation on the structure of other domains. Accordingly, the protein of the present invention may or may not comprise a linker domain. The protein may alternatively comprise a portion of a linker domain, and it is preferable if the protein retains a fragment of a linker domain comprising about 10 amino acid residues.

[0024] The other aspect of the present invention relates to a protein comprising any of the amino acid sequences as shown in SEQ ID NO: 1 (RCE I), SEQ ID NO: 3 (RCE II), SEQ ID NO: 5 (RCE III), SEQ ID NO: 7 (MCE I), SEQ ID NO: 9 (MCE II), or SEQ ID NO: 11 (PCE I), wherein the cellulose-binding domain has been deleted therefrom, and exhibiting endoglucanase activity. The present invention further relates to a modified protein and a homologue of such protein exhibiting endoglucanase activity.

[0025] In this description, the term "modified protein" refers to a protein that comprises an amino acid sequence having modification such as addition, insertion, diminution, deletion, or substitution of one or several amino acids in the amino acid sequence of RCE I, RCE II, RCE III, MCE I, MCE II, or PCE I, which lacks the cellulose-binding domain. The number of the amino acids to be involved with such modification is not particularly limited as long as the modified

protein has endoglucanase activity. The number thereof is preferably 1 to about 50, more preferably 1 to about 30, and still more preferably 1 to 9.

[0026] The term "homologue" used herein refers to a polypeptide having an amino acid sequence coded by a gene encoding the amino acid sequences of RCE I, RCE II, RCE III, MCE I, MCE II, and PCE I lacking the cellulose-binding domain. One example would be a gene (nucleotide sequence) that hybridizes under stringent conditions with DNA having any nucleotide sequence as shown in SEQ ID NO: 2 or 13 (RCE I), SEQ ID NO: 4 (RCE II), SEQ ID NO: 6 (RCE III), SEQ ID NO: 8 (MCE I), SEQ ID NO: 10 (MCE II), or SEQ ID NO: 12 (PCE I), wherein a portion encoding a cellulose-binding domain has been removed and having endoglucanase activity. The term "stringent conditions" refers to conditions under which, while a probe that comprises the nucleotide sequence encoding a part or all of the amino acid sequences of RCE I, RCE II, RCE III, MCE I, MCE II, and PCE I lacking the cellulose-binding domain or an amino acid sequence of its modified protein hybridizes with a gene encoding a homologue, this probe is controlled to such an extent that it does not hybridize with the endoglucanase NCE4 gene (SEQ ID NO: 18) according to WO 98/03640 or the endoglucanase SCE3 gene (SEQ ID NO: 19) according to WO 98/54332 (wherein the amount of DNA used is equal to that of the gene encoding the NCE4 gene, SCE3 gene, or a homologue). A specific example of "stringent conditions" is as follows. A labeled probe having a full length DNA sequence encoding amino acid sequences such as RCE I, which lacks the cellulose-binding domain, is used. In accordance with the method of the ECL Direct DNA/RNA Labeling Detection System (Amersham), prehybridization is carried out at 42°C for 1 hour, the probe is added, and hybridization is then carried out at 42°C for 15 hours. Thereafter, 0.5x SSC (1x SSC; 15 mM trisodium citrate, 150 mM sodium chloride) comprising 0.4% SDS and 6M urea is used to perform washing twice at 42°C for 20 minutes. Subsequently, 5x SSC is used to perform washing twice at room temperature (about 25°C) for 10 minutes.

[0027] Examples of such modified proteins or homologue include a protein having an amino acid sequence that is preferably at least 70%, more preferably at least 80%, still more preferably at least 90%, even more preferably at least 95%, and most preferably at least 98% homologous to the amino acid sequence of RCE I, RCE II, RCE III, MCE I, MCE II, or PCE I, which lacks the cellulose-binding domain. The numerical values indicating homology may be determined using a program for searching for homology that is known to a person skilled in the art. Preferably, these numerical values are determined using default (initial setting) parameters at FASTA3 (Science, 227, 1435-1441 (1985); Proc. Natl. Acad. Sci. USA, 85, 2444-2448 (1988); <http://www.ddbj.nig.ac.jp/E-mail/homology-j.html>).

[0028] The protein of the present invention does not comprise a cellulose-binding domain. Accordingly, the modified protein and the homologue should not comprise a cellulose-binding domain. This can be confirmed by investigating the amino acid sequence of the object protein and whether or not the consensus sequence (Hoffren, A. -M. et al., Protein Engineering 8: 443-450, 1995; SEQ ID NO: 17) is present therein.

[0029] Also, the protein of the present invention has endoglucanase activity. Accordingly, the modified protein and the homologue should also have endoglucanase activity. This can be confirmed by investigating endoglucanase activity of the object protein by the aforementioned method.

[0030] The protein of the present invention can be produced as a protein comprising an amino acid sequence that does not comprise a cellulose-binding domain by a method known to a person skilled in the art based on the amino acid sequence of a known endoglucanase derived from *Zygomycetes*. Examples of such a method include a method that is carried out by decomposing in a linker domain using protease during the culture of *Zygomycetes* producing endoglucanase to cause deletion of the cellulose-binding domain and a method that is carried out by artificially expressing endoglucanase that does not have a cellulose-binding domain due to genetic engineering techniques. Among the proteins of the present invention, in particular, the modified protein and the homologue can be prepared using DNA encoding their amino acid sequences by a genetic engineering technique that is known to a person skilled in the art.

[0031] The protein of the present invention can yield a higher effect than the original endoglucanase having a cellulose-binding domain in fabric treatment or other applications to detergents or paper pulp. Particularly, a much higher effect can be attained in the activity of removing fuzz from reproduced cellulose fabric such as lyocell (per protein weight) and in the activity of removing fuzz from cotton fabric such as knitted cotton (per protein weight). The protein of the present invention is preferably twice or higher, more preferably 2.5 times or higher, and most preferably 3 times or higher in the activity of removing fuzz from reproduced cellulose fabric (such as lyocell) (per protein weight) as a purified endoglucanase having the cellulose-binding domain. Or, the protein is preferably 5 times or higher, more preferably 15 times or higher, and most preferably 20 times or higher in the activity of removing fuzz from cotton fabric such as knitted cotton (per protein weight) as a purified endoglucanase having the cellulose-binding domain.

2. Gene, expression vector, host cell transformed with the expression vector, and production of endoglucanase lacking the cellulose-binding domain using the host cell

[0032] The present invention relates to a gene encoding the protein of the present invention, a modified protein thereof, or a homologue of the protein or the modified protein and an expression vector comprising the gene.

[0033] The gene of the present invention may be any gene as long as it encodes the protein of the present invention,

a modified protein thereof, or a homologue of the protein or the modified protein. Specific details of their nucleotide sequences are not particularly limited. Examples of usable genes in order to express RCE I, RCE II, RCE III, MCE I, MCE II, or PCE I, which lacks the cellulose-binding domain, as the protein of the present invention include those comprising nucleotide sequences as shown in SEQ ID NO: 2 (RCE I), SEQ ID NO: 4 (RCE II, SEQ ID NO: 6 (RCE III),
 5 SEQ ID NO: 8 (MCE I), SEQ ID NO: 10 (MCE II), or SEQ ID NO: 12 (PCE I).

[0034] Specifically, the gene of the present invention comprises an amino acid sequence as shown in SEQ ID NO: 1, 3, 5, 7, 9, or 11, wherein the cellulose-binding domain has been deleted therefrom, and encodes a protein exhibiting endoglucanase activity, a modified protein thereof exhibiting endoglucanase activity, or a homologue of the protein or the modified protein exhibiting endoglucanase activity.

10 **[0035]** The gene of the present invention comprises DNA as described in the following (a) or (b).

(a) DNA that has a nucleotide sequence as shown in SEQ ID NO: 2 or 13, 4, 6, 8, 10, or 12, wherein a portion encoding a cellulose-binding domain has been deleted therefrom.

15 (b) DNA that hybridizes under stringent conditions with DNA that has a nucleotide sequence as shown in SEQ ID NO: 2 or 13, 4, 6, 8, 10, or 12, wherein a portion encoding a cellulose-binding domain has been deleted therefrom.

[0036] The term "stringent conditions" means the conditions described in the section 1. above.

[0037] The nucleotide sequences of the aforementioned genes can be optimized depending on the type of host cell used in the later transformation. These nucleotide sequences can be optimized with respect to, for example, the codon usage in a host cell or the intron recognition sequence in a host cell. The codon usage can be optimized by, for example,
 20 modifying nucleotide sequences so as to comprise as many codons used in a host cell at high frequency as possible without changing the amino acid sequence to be coded. This can improve the efficiency of translation from genes into proteins. The intron recognition sequence can be optimized by, for example, modifying nucleotide sequences so as to have no DNA sequence, which could be recognized as an intron in a host cell, or to have as few sequences as possible
 25 without changing the amino acid sequence to be coded. This can improve the stability of mRNA that is a transcript of an object gene. The intron recognition sequences vary depending on types of host cells. Examples of intron recognition sequences of filamentous fungi belonging to the Fungi Imperfecti include DNA sequences such as GTAGN, GTATN, GTAAN, GTACGN, GTGTN, GCACGN, and GTTCGN ("N" stands for A, T, C, or G in each sequence).

[0038] In this description, the term "codon optimized gene" refers to a gene that is obtained by optimizing the codon usage and/or the intron recognition sequence. Preferably, the codon optimized gene is a gene obtained by optimizing the codon usage, and more preferably a gene obtained by optimizing the codon usage and the intron recognition sequence. This codon optimized gene is preferably optimized for the expression in filamentous fungi belonging to the Fungi imperfecti. Examples of such codon optimized genes include a gene which lacks a portion encoding a cellulose-binding domain in the codon optimized endoglucanase RCE I gene (SEQ ID NO: 13) as disclosed in WO 00/24879.

35 **[0039]** The expression vector of the present invention comprises an object gene (a gene encoding the protein of the present invention, a modified protein thereof, or a homologue of the protein or the modified protein) in a state that is replicable in a host cell and expressible of the protein of the present invention, a modified protein thereof, or a homologue of the protein or the modified protein. This expression vector can be constructed based on a self-replicating vector, i. e., a vector that exists as an extrachromosomal entity and replicates independently of the chromosome, for example,
 40 a plasmid. Alternatively, the expression vector may be a vector that is integrated into the genome of the host cell upon introduction therein and replicated together with the chromosome into which it has been incorporated. For the construction of the vector of the present invention, conventional procedures and methods used in the field of genetic engineering can be used.

[0040] For the expression of the protein of the present invention, a modified protein thereof, or a homologue of the protein or the modified protein upon introduction into the host cell, it is desirable that the expression vector of the invention contains DNA sequences to regulate the expression and gene markers, etc. to select transformants, in addition to the gene encoding the protein of the present invention, a modified protein thereof, or a homologue of the protein or the modified protein. Examples of expression regulatory DNA sequences include promoters, terminators,
 45 and DNA sequences encoding signal peptides. The promoters and the terminators are not particularly limited as long as they show transcription activity in the host cell. They may be obtained as DNA sequences, which control the expression of a gene encoding a protein homogeneous or heterogeneous to the host cell. The signal peptides are not particularly limited as long as they contribute to the secretion of protein in the host cell. They may be obtained from DNA sequences derived from a gene encoding a protein homogeneous or heterogeneous to the host cell. The gene markers of the invention can be appropriately selected depending on the method for selecting transformants. For
 50 example, genes encoding drug resistance or genes complementing auxotrophy may be used. Each of these DNA sequences and gene markers is operably linked to the expression vector of the present invention.

55 **[0041]** Further, the present invention relates to a host cell that is transformed with the expression vector. The expression vector introduced into a host cell should be replicable therein. Thus, the host cell that is used herein varies

depending on the type of vector used in the production of the expression vector. Alternatively, in accordance with the type of a host cell to be used, an expression vector can be produced so as to be replicable therein. Specifically, in order to obtain a transformant that expresses the protein of the present invention, a modified protein thereof, or a homologue of the protein or the modified protein, a host cell and an expression vector should be adequately combined. Such a combination is referred to as a host-vector system. The host-vector system that is used in the present invention is not particularly limited. Examples thereof include systems using microorganisms such as *Escherichia coli*, *Actinomyces*, yeast, and filamentous fungi as host cells, and a system using filamentous fungi is preferred. An expression system for a fusion protein with other protein can be also used.

[0042] When filamentous fungi are used as host cells, any type of filamentous fungi can be used, and preferred examples thereof include those belonging to *Humicola*, *Aspergillus*, or *Trichoderma*. Particularly preferred examples of these filamentous fungi include *Humicola insolens*, *Aspergillus niger* or *Aspergillus oryzae*, and *Trichoderma viride*.

[0043] A host cell can be transformed with the expression vector of the present invention in accordance with conventional methods used in the field of genetic engineering.

[0044] The thus obtained transformant (transformed host cell) is cultured in a suitable medium, and the protein of the present invention, a modified protein thereof, or a homologue of the protein or the modified protein can be isolated and obtained from the culture product. Accordingly, another aspect of the present invention relates to a method for producing a protein comprising steps of culturing the host cell of the present invention and collecting the protein of the present invention, a modified protein thereof, or a homologue of the protein or the modified protein from the host cell obtained by the step of culture or a culture product thereof. Culture methods and other conditions for transformants may be substantially the same as those for microorganisms to be used. The transformants can be cultured and the object protein can be then collected by a conventional method of this technical field.

3. Cellulase preparation

[0045] The present invention relates to a cellulase preparation that comprises the protein of the present invention, a modified protein thereof, a homologue of the protein or the modified protein, or a protein produced by the method for producing a protein according to the present invention.

[0046] In general, a cellulase preparation is powder, liquid, or the like that comprises, for example, an excipient (e.g., lactose, sodium chloride, or sorbitol), a preservative, or a nonionic surfactant, in addition to a cellulase enzyme. For example, it is formulated as a powdery, particulate, granular, non-dusting granular, or liquid preparation. The cellulase preparation of the present invention comprises, as the cellulase enzyme, the protein of the present invention, a modified protein thereof, a homologue of the protein or the modified protein, or a protein produced by the method for producing a protein of the present invention (hereinafter referred to as "the proteins of the present invention"). Further, the cellulase preparation of the present invention may comprise, in addition to the proteins of the present invention, other cellulase enzymes, for example, cellobiohydrolase, β -glucosidase, or endoglucanase which are not involved in the present invention.

[0047] One type of cellulase preparation, a non-dusting granular preparation, may be prepared by a conventional dry granulating method. Specifically, the powdery proteins of the present invention are mixed with one or several of: neutral inorganic salts that do not affect endoglucanase activity represented by sodium sulfate, sodium chloride, or the like; minerals that do not affect endoglucanase activity represented by bentonite, montmorillonite, or the like; or neutral organic substances represented by starch, particulate cellulose, or the like. A powder or fine suspension of one or several nonionic surfactant is then added thereto, followed by thorough mixing or kneading. Depending on the situation, a synthetic polymer represented by polyethylene glycol for binding solid matter or a natural polymer such as starch is suitably added and further kneaded. Thereafter, extrusion granulation is carried out using, for example, Disc Pelleter, and the extruded granules are then shaped into spherical form using the Marumerizer, followed by drying. Thus, non-dusting granules can be produced. The amount of one or several nonionic surfactants to be added is not particularly limited. The amount is preferably 0.1% to 50% by weight, more preferably 0.1% to 30% by weight, and still more preferably 1% to 20% by weight, based on the entire cellulase preparation according to the present invention. Oxygen permeation or water permeation can be regulated by coating the surfaces of granules with a polymer, etc.

[0048] In contrast, a liquid formulation can be prepared by incorporating a stabilizer for the endoglucanase enzyme such as a synthetic polymer, a natural polymer, or the like into a solution comprising the proteins of the present invention and adding inorganic salts or synthetic preservatives according to need. In this case, one or several nonionic surfactants can be also incorporated. The amount of one or several nonionic surfactants to be added is not particularly limited. The amount is preferably 0.1% to 50% by weight, more preferably 0.1% to 30% by weight, and still more preferably 1% to 20% by weight, based on the entire cellulase preparation according to the present invention.

4. Application of the proteins of the present invention and the cellulase preparation of the present invention

[0049] The present invention relates to a method for treating cellulose-containing fabrics. This method comprises a step of bringing cellulose-containing fabrics into contact with the proteins of the present invention or the cellulase preparation of the present invention. Conditions such as contact temperature or amounts of the proteins or the cellulase preparation can be suitably determined with respect to various other conditions.

[0050] The aforementioned method can be used to reduce the rate at which cellulose-containing fabrics become fuzzy or to reduce fuzzing in cellulose-containing fabrics. In this application, the proteins or the cellulase preparation at the protein concentration of 0.001 to 1 mg/l is preferably used at about 30°C to 60°C.

[0051] The aforementioned method can be used in weight loss treatment of cellulose-containing fabrics to improve the feel and appearance thereof. In this application, to improve the feel means to reduce the rate at which the feel is spoiled. In this application, the proteins or the cellulase preparation at the protein concentration of 0.001 to 100 mg/l is preferably used at about 30°C to 60°C.

[0052] The aforementioned method can be used for color clarification of colored cellulose-containing fabrics.

[0053] The aforementioned method can be used for providing colored cellulose-containing fabrics with partial color change. In this application, for example, colored cellulose-containing fabrics (e.g., denim) can be provided with an appearance of stone-washed material. In this application, the proteins or the cellulase preparation at the protein concentration of 0.01 to 100 mg/l is preferably used at about 40°C to 60°C.

[0054] The protein concentrations of various endoglucanases were calculated from the peak area at UV 280 nm of respective endoglucanase eluted with a linear gradient from 0% to 80% of acetonitrile concentration in 0.05% TFA (trifluoroacetic acid) at a flow rate of 1.0 ml/min in HPLC analysis using TSK gel TMS-250 column (4.6 mm I.D. x 7.5 cm, TOSOH Japan). The standard used was the purified NCE4, which was analyzed in HPLC under the same conditions, the protein concentration of which had been preliminarily measured by a protein assay kit (BioRad Laboratories). The purified NCE4 is purified from a culture product of *Humicola insolens* MN 200-1, which was deposited at the International Patent Organism Depositary of the National Institute of Advanced Industrial Science and Technology (Tsukuba Central 6, 1-1-1 Higashi, Tsukuba, Ibaraki, Japan) under the accession number of FERM BP-5977 (initial deposition: FERM P-15736, date of initial deposition: July 15, 1996), in accordance with the method as described in WO 98/03640. The standard used to measure the protein concentration using the protein assay kit is the Albumin Standard (Bovine serum albumin, fraction V, PIERCE).

[0055] The aforementioned method can be used to reduce the rate at which cellulose-containing fabrics become stiff or to reduce stiffness in cellulose-containing fabrics. In this application, cellulose-containing fabrics can be softened.

[0056] In the aforementioned applications, cellulose-containing fabrics can be treated through soaking, washing, or rinsing of the fabrics. Specifically, the aforementioned method of the present invention can be carried out by treating cellulose-containing fabrics during washing. On some occasions, however, the treatment of fabrics may be carried out during soaking or rinsing by adding the proteins or the cellulase preparation of the present invention into water where the fabrics are soaked or to be soaked.

[0057] The present invention relates to an additive to detergent comprising the proteins or the cellulase preparation of the present invention in a non-dusting granular form or a stabilized liquid form. The present invention further relates to a detergent composition comprising the proteins or the cellulase preparation of the present invention.

[0058] This detergent composition may also contain a surfactant (which may be anionic, nonionic, cationic, amphoteric, or zwitterionic surfactant, or a mixture thereof). Further, this detergent composition may contain other detergent components known in the art, such as builders, bleaching agents, bleaching activators, corrosion inhibitors, sequestering agents, soil-dissociating polymers, aromatics, other enzymes (e.g., protease, lipase, or amylase), enzyme stabilizers, formulation assistants, fluorescent brightening agents, foaming promoters, etc. Examples of representative anionic surfactants include linear alkyl benzene sulfonate (LAS), alkylsulfate (AS), α -olefin sulfonate (AOS), polyoxyethylene alkyl ether sulfate (AES), α -sulfonato fatty acid methyl ester (α -SFMe), and alkali metal salts of natural fatty acids. Examples of nonionic surfactants include polyoxyethylene alkyl ether (AE), alkyl polyethylene glycol ether, nonylphenol polyethylene glycol ether, fatty acid methyl ester ethoxylate, fatty acid esters of sucrose or glucose, and esters of alkyl glucoside and polyethoxylated alkylglucoside.

[0059] The use of the proteins or the cellulase preparation of the present invention in a detergent composition can improve performances regarding particulate soil removal, color clarification, fuzz prevention, depilling, and reduction of stiffness.

[0060] The present invention relates to a method of deinking waste paper using a deinking agent wherein the proteins or the cellulase preparation of the present invention is used in a step of deinking waste paper with a deinking agent.

[0061] When the protein or the cellulase preparation of the present invention is acted on waste paper, the efficiency of deinking is enhanced, and thus, the protein or the cellulase preparation of the present invention is useful in the process of manufacturing recycled paper from waste paper. This deinking method can significantly decrease ink-remaining fibers. Therefore, the whiteness of the waste paper can be enhanced.

[0062] The aforementioned "deinking agent" is not particularly limited as long as it is a commonly used agent when deinking waste paper. Examples thereof include alkali such as NaOH or Na₂CO₃, sodium silicate, hydrogen peroxide, phosphates, anionic surfactants, nonionic surfactants, capturing agents such as oleic acid, and examples of aids include pH stabilizers, chelating agents, and dispersants.

[0063] Waste paper, which can be treated by the above deinking method, is not particularly limited as long as it can be generally referred to as waste paper. Examples of waste paper include: used printed paper containing mechanical pulp and chemical pulp such as used newspaper, used magazine paper and low-grade or middle-grade used printed paper, used wood-free paper composed of chemical pulp; and coated paper thereof. Further, the above deinking method can be applied to any paper on which ink has been deposited if the paper is not generally referred to as waste paper.

[0064] Further, the present invention relates to a method for improving drainage of paper pulp. This method comprises a step of treating the paper pulp with the proteins or the cellulase preparation of the present invention.

[0065] According to this method, drainage of paper pulp can be remarkably improved without significant deterioration in paper strength. Pulp, which can be treated by this method, is not particularly limited, and examples thereof include waste paper pulp, recycled board pulp, kraft pulp, sulfite pulp, processed or thermo-mechanical pulp, and other high-yield pulp.

[0066] The present invention further relates to a method for improving the digestibility of animal feeds. This method comprises a step of treating animal feeds with the proteins or the cellulase preparation of the present invention.

[0067] According to this method, molecular weights of glucans in animal feeds are suitably lowered. Thus, the digestibility of animal feeds can be improved.

5. Deposition of microorganisms

[0068] The *Rhizopus oryzae* CP96001 strain was deposited at the International Patent Organism Depository of the National Institute of Advanced Industrial Science and Technology (Tsukuba Central 6, 1-1-1 Higashi, Tsukuba, Ibaraki, Japan) under the accession number of FERM BP-6889 on April 21, 1997.

[0069] The *Mucor circinelloides* CP99001 strain was deposited at the International Patent Organism Depository of the National Institute of Advanced Industrial Science and Technology (Tsukuba Central 6, 1-1-1 Higashi, Tsukuba, Ibaraki, Japan) under the accession number of FERM BP-6890 on July 2, 1999.

[0070] The *Phycomyces nitens* CP99002 strain was deposited at the International Patent Organism Depository of the National Institute of Advanced Industrial Science and Technology (Tsukuba Central 6, 1-1-1 Higashi, Tsukuba, Ibaraki, Japan) under the accession number of FERM BP-6891 on July 2, 1999.

[0071] The *Escherichia coli* JM 109 strain that was transformed with the expression vector pMKD01 used in the present invention was deposited at the International Patent Organism Depository of the National Institute of Advanced Industrial Science and Technology (Tsukuba Central 6, 1-1-1 Higashi, Tsukuba, Ibaraki, Japan) under the accession number of FERM BP-5974 (initial deposition: PERM P-15730, date of initial deposition: July 12, 1996).

[0072] The *Humicola insolens* MN 200-1 strain that can be a host for the expression vector of the present invention was deposited at the International Patent Organism Depository of the National Institute of Advanced Industrial Science and Technology (Tsukuba Central 6, 1-1-1 Higashi, Tsukuba, Ibaraki, Japan) under the accession number of FERM BP-5977 (initial deposition: FERM P-15736, date of initial deposition: July 15, 1996).

[0073] This description includes part or all of the contents as disclosed in the description of Japanese Patent Application No. 2000-354296, which is a priority document of the present application.

Best Modes for Carrying out the Present Invention

[0074] The present invention is described in more detail with reference to the following examples and reference examples but is not limited thereto.

[0075] In the following description, "endoglucanase activity" refers to CMCase activity. Further, "CMCase activity" measured by the amount of the reducing sugars released after the incubation of a cellulase enzyme and a carboxymethylcellulose (CMC, Tokyo Kasei Kogyo, Japan) solution for a given period of time and one unit is defined as the amount of an enzyme that produces reducing sugars corresponding to 1 μ mol of glucose per minute.

[0076] In implementing the following tests, examples disclosed in WO 98/03667 and WO 00/24879 were referred to.

[Reference Examples]

Reference Example 1: cDNA cloning of cellulase NCE5

5 (1) Isolation of cDNA and preparation of library

[0077] In the screening of the gene that encodes NCE5, the cellulase component, mRNA was prepared from *Humicola insolens* MN 200-1 (FERM BP-5977), and cDNA was synthesized by a reverse transcriptase to prepare a library.

10 (i) Preparation of total RNA

[0078] *Humicola insolens* MN 200-1 (FERM BP-5977) was cultured in medium (N) (5.0% Avicel, 2.0% yeast extract, 0.1% polypeptone, 0.03% calcium chloride, 0.03% magnesium chloride, pH 6.8) for 2 days, and cells were collected by centrifugation (3,500 rpm, 10 minutes). Among the collected cells, 3 g thereof was washed with sterilized water, frozen in liquid nitrogen, and then ground in liquid nitrogen using a mortar and a pestle. Total RNA was isolated from the ground cells using ISOGEN (Nippon Gene) in accordance with the manual attached thereto, and total RNA was confirmed by formaldehyde agarose gel electrophoresis as a chromatic figure.

20 (ii) Preparation of poly(A)tail+RNA (= mRNA)

[0079] Among the total RNA prepared in (i), 1 mg thereof was applied on the oligo (dT) cellulose column to elute and isolate mRNA using the mRNA Purification Kit (Amersham Pharmacia Biotech) in accordance with the attached manual. Further, mRNA was confirmed by formaldehyde agarose gel electrophoresis as a smear chromatic figure.

25 (iii) Synthesis of cDNA

[0080] cDNA was synthesized from 5 µg of the mRNA prepared in (ii) using the Time Saver cDNA Synthesis Kit (Amersham Pharmacia Biotech) in accordance with the attached manual.

30 (iv) Preparation of cDNA library

[0081] The *Eco*RI-NotI adaptor contained in the Time Saver cDNA Synthesis Kit was ligated to the blunt end of the synthesized total cDNA in accordance with the attached manual. The total amount of this DNA fragment was ligated into the phage vector and the *Eco*RI arm of the λZAPII Cloning Kit (Stratagene) using the DNA Ligation Kit Ver. 2 (Takara Shuzo Co., Ltd.), followed by ethanol precipitation. Thereafter, the DNA fragment was dissolved in a TE (10 mM Tris hydrochloride, pH 8.0, 1 mM EDTA) buffer. The thus obtained recombinant phage vector was subjected to *in vitro* packaging using the Gigapack III Plus Packaging Extract (Stratagene) in accordance with the attached manual. Thereafter, this recombinant phage vector was infected with *Escherichia coli* XL1-Blue MRF' and cultured on a plate for plaque formation. Thus, a phage library was obtained. With the use thereof, an object gene was cloned.

40 (2) Amplification and analysis of DNA by PCR

[0082] DNA was amplified by PCR using the cDNA prepared in (i)-(iii) as a template based on the information concerning partial amino acid sequences of the cellulase NCE5.

[0083] The following synthetic oligonucleotides were prepared as primers.
N-terminal: 5'-TAY TGG GAY TGY TGY AAR CC-3' (20mer) (SEQ ID NO: 36)
T-43.0: 5'-TCI GCR TTI ARR AAC CAR TC-3' (20mer) (SEQ ID NO: 37)

(In these nucleotide sequences, R indicates G or A; Y indicates T or C; and I indicates inosine.)

[0084] PCR was carried out in 50 µl of reaction solution using 1 µg of cDNA as a template, 1.25 units of LA Taq DNA Polymerase (Takara Shuzo Co., Ltd.) and its attached buffer, 0.2 mM dNTP, 10% DMSO, and 1 µM each of the above primers under the following condition: at 94°C for 1 minute, (at 94.0°C for 30 seconds, 55.0°C for 30 seconds, and 72.0°C for 1 minute) x 25 times, and 72.0°C for 5 minutes.

[0085] About 500 bp DNA was amplified by this reaction, and the amplified DNA was subjected to sequencing using the DYEnamic ET Terminator Cycle Sequencing Premix Kit (Amersham Pharmacia Biotech) and the ABI PRISM 310 Genetic Analyzer (PE Applied Biosystems) in accordance with the attached protocols. As a result, the amino acid sequence, which was deduced from the determined nucleotide sequence, contained all the partial amino acid sequence-

es of cellulase NCE5. Thus, the deduced amino acid sequence was used as a probe in the following screening process.

(3) Cloning of the gene that encodes NCE5, the cellulase component

5 (i) Screening by plaque hybridization

[0086] The 500 bp DNA fragment (100 ng) amplified by PCR was previously labeled with the ECL Direct DNA/RNA Labeling Detection System (Amersham Pharmacia Biotech).

10 [0087] The phage plaque prepared in (1)-(iv) was transferred to the Hybond-N+ nylon transfer membrane (Amersham Pharmacia Biotech) and subjected to alkali treatment with 0.4N sodium hydroxide. Recombinant phage DNA on the membrane was denatured into a single strand. Thereafter, the nylon membrane was washed with 5x SSC (1x SSC: 15 mM trisodium citrate, 150 mM sodium chloride) and then air dried to immobilize DNA thereon. Thereafter, the nylon membrane was hybridized in accordance with the manual of the kit, detected, and exposed to Fuji Medical X-ray film (Fuji Photo Film) to yield 6 positive clones.

15 (ii) Preparation of phage DNA

[0088] DNA was prepared from the positive clones as plasmid DNA in accordance with the manual attached to the kit.

20 [0089] A plasmid in which a DNA fragment was cloned into pBluescript SK(-) was prepared from ampicillin-resistant *Escherichia coli* SOLR™. Using this plasmid as a template, PCR was carried out using the N-terminal and T-43.0 primers used in (2) under the same conditions as described above. As a result, a 500 bp amplification product was obtained from one plasmid. Thus, it is deduced that the object DNA was cloned into this plasmid. This plasmid was digested with *EcoRI* and then subjected to agarose gel electrophoresis.

25 [0090] As a result, the plasmid was found to contain about 1 kbp *EcoRI* fragment.

(4) Determination of cDNA nucleotide sequence

30 [0091] A nucleotide sequence of the about 1 kbp *EcoRI* fragment that was inserted into a positive recombinant pBluescript SK(-) plasmid obtained in (3)-(ii) was determined in the same manner as described above using primers for sequencing T3 and T7. As a result, this nucleotide sequence was found to contain a 672 bp open reading frame (ORF). Amino acid sequences deduced from the nucleotide sequence and the ORF were shown in SEQ ID NO: 39 and SEQ ID NO: 38 in the Sequence Listings.

35 [0092] Further, 1 to 18 amino acid sequences in this ORF were considered to be signal sequences for secreting the protein extracellularly.

Reference Example 2: Expression of NCE5 gene in *Humicola insolens*

40 [0093] Plasmid pJD01 (see Example D1 (2) (b) in WO 00/24879) was used as an expression vector in *Humicola insolens* MN 200-1 (PERM BP-5977) and constructed in the following manner.

(1) Construction of NCE5 expressing plasmid pJND-c5

(i) Site-directed mutagenesis into NCE5 gene

45 [0094] In order to ligate the NCE5 gene to the *Bam*HI site of plasmid pJD01, a primer was constructed so as to previously comprise *Bam*HI site in a sequence immediately upstream of the initiation codon and that immediately downstream of the termination codon, and the NCE5 gene was amplified by PCR. Primers for mutagenesis were designed as shown below.

NCE5-N-*Bam*HI:

50

5'-GGGGATCCTGGGACAAGATGCAGCTCCCCCTGACCACG-3' (38mer) (SEQ ID NO: 40);

55 NCE5-C-*Bam*HI:

5'-GGGGATCCTGCATTAAACGCGAGCAGCCGCTCTTGGCC-3' (38mer) (SEQ ID NO: 41).

[0095] PCR was carried out under the same conditions as described above using the positive recombinant pBlue-script SK(-) plasmid obtained in Reference Example 1 as a template. As a result, an about 670 bp amplification product of the DNA fragment was confirmed by 1.0% agarose gel electrophoresis. Unreacted matter was removed by the MicroSpin S-400 HR Columns (Amersham Pharmacia Biotech), precipitated with ethanol, and then digested with *Bam*-HI. Subsequently, a total amount was subjected to 1.0% agarose gel electrophoresis, a 670 bp DNA fragment was collected using the Sephaglas BandPrep Kit (Amersham Pharmacia Biotech) in accordance with the attached manual, and the *Bam*HI fragment was subcloned into the *Bam*HI site of plasmid pUC118 to yield plasmid pNCE5Bam. Further, the nucleotide sequence of this inserted fragment was determined and confirmed by the aforementioned method.

(ii) Preparation of plasmid pJND-c5

[0096] The above plasmid pJD01 was digested with *Bam*HI and separated by 0.8% agarose gel electrophoresis. An about 8.0 kbp DNA fragment was collected by the Sephaglas BandPrep Kit, and the collected DNA fragment was dephosphorylated using *Escherichia coli*-derived alkaline phosphatase (Takara Shuzo Co., Ltd.) in accordance with the attached manual. Similarly, the plasmid pNCE5Bam obtained in (i) was digested with *Bam*HI, and 670 bp DNA fragments were collected and ligated to the above DNA fragment using the DNA Ligation Kit Ver. 2 to obtain expression plasmid pJND-c5.

(2) Transformation of *Humicola insolens* with plasmid pJND-c5

[0097] *Humicola insolens* MN 200-1 (FERM BP-5977) was cultured in medium (S) at 37°C for 24 hours, and then the cells were collected by centrifugation at 3,000 rpm for 10 minutes. Medium (S) is composed of medium (N) described in Reference Example 1 having glucose (3.0%) added thereto and Avicel removed therefrom. The collected cells were washed with 0.5M sucrose and suspended in 10 ml of enzyme solution for preparing protoplast (3 mg/ml β -glucuronidase, 1 mg/ml Chitinase, 1 mg/ml Zymolyase, and 0.5M sucrose) filtered through a 0.45 μ m filter. The suspension was shaken at 30°C for 60 to 90 minutes to render fungal threads to be protoplasted. This suspension was filtered and then centrifuged at 2,500 rpm for 10 minutes, and the protoplast was collected and then washed with a SUTC buffer (0.5M sucrose, 10 mM calcium chloride, and 10 mM Tris hydrochloride (pH 7.5)).

[0098] The thus prepared protoplast was suspended in 1 mL of SUTC buffer, and 10 μ l of DNA (TE) solution was added to the suspension in amounts of 10 μ g per each 100 μ l of the suspension. The mixture was allowed to stand in ice for 5 minutes. Subsequently, 400 μ l of PEG solution (60% PEG 4000, 10 mM calcium chloride, 10 mM Tris hydrochloride (pH 7.5)) was added and the mixture was allowed to stand in ice for 20 minutes. Thereafter, 10 ml of SUTC buffer was added, and centrifugation was carried out at 2,500 rpm for 10 minutes. The collected protoplast was suspended in 1 ml of SUTC buffer, centrifuged at 4,000 rpm for 5 minutes, and finally suspended in 100 μ l of SUTC buffer.

[0099] The protoplast treated as above was superposed on the hygromycin-added (200 μ g/ml) YMG medium (1% glucose, 0.4% yeast extract, 0.2% malt extract, 1% agar (pH 6.8)) together with YMG soft agar. Culture was incubated at 37°C for 5 days. Thereafter, the generated colony was determined to be a transformant.

[Example 1] Construction of gene expressing RCE I variant H43, which lacks the cellulose-binding domain

[0100] Plasmid p18-1 containing the codon optimized endoglucanase gene RCE I (see Example D3 (1) g) in WO 00/24879) was digested with the restriction enzyme *Bam*HI to prepare plasmid pR1H4 wherein a fragment containing the codon optimized endoglucanase gene has been cloned into the *Bam*HI site of plasmid pUC118. This plasmid pR1H4 was used as a template to perform first-phase PCR using two synthetic DNAs, i.e., RC-43F and RC-43R, as primers and the TaKaRa LA PCR in vitro Mutagenesis Kit (Takara Shuzo Co., Ltd.). Reaction conditions were in accordance with the manual attached to the kit. A sample was separated after the reaction by agarose gel electrophoresis to obtain an about 650 bp gene fragment 43-X2.

[0101] Using plasmid pJND-c5 as a template, which comprises the NCE5 gene originally having no cellulose-binding domain and being a family 45 endoglucanase as described in Reference Example 2, first-phase PCR was similarly carried out using two synthetic DNAs, i.e., NX-43F and NX-43R, as primers and the TaKaRa LA PCR in vitro Mutagenesis Kit (Takara Shuzo Co., Ltd.). Reaction conditions were in accordance with the manual attached to the kit. A sample was separated after the reaction by agarose gel electrophoresis to obtain an about 120 bp gene fragment 43-X1.

RC-43F: CACCACGCGCTACTGGGACT (SEQ ID NO: 20);

RC-43R: GGATCCTGCGTTTACTTGC (SEQ ID NO: 21);

NX-43F: GGATCCTGGGACAAGATG (SEQ ID NO: 22);

NX-43R: GCACGACGGCTTGCAGC (SEQ ID NO: 23)

[0102] Annealing and second-phase PCR were carried out using PCR fragments 43-X1 and 43-X2 and the TaKaRa LA PCR in vitro Mutagenesis Kit. Two synthetic DNAs, i.e., NX-43F and RC-43R, were used as primers and reaction conditions were in accordance with the manual attached to the kit. A sample was separated after the reaction by agarose gel electrophoresis to obtain an about 700 bp gene fragment H43. This fragment was digested with the restriction enzyme *Bam*HI to prepare plasmid pR1H43 ligated to the *Bam*HI site of plasmid pUC118 using the TaKaRa DNA Ligation Kit Ver. 1. Reaction conditions, such as those regarding enzymes, were in accordance with the manual attached to the kit. The obtained plasmid pR1H43 was subjected to sequencing using the Cy5-Auto Read Sequencing Kit (Amersham Pharmacia), and the sequence was analyzed using the DNA Sequencer ALFred (Amersham Pharmacia). A primer used in the reaction was the M13 primer, which was attached to the kit. As a result, it was confirmed that the sequence was in the expected form, i.e., a secretion signal on the N-terminal side was derived from NCE5 and the remaining catalytic active domain (CAD) was a sequence derived from RCE I. The amino acid sequence of the RCE I variant H43 deduced from the nucleotide sequence is shown below.

H43: MQLPLTLLTLLPALAAQSGSGRTRYWDCKPSCSWPGKANVSSPVKSCNKDG
VTALSDSNAQSGCNGGNSYMCNDNQPWAVNDNLAYGFAAAISGGGESRWCCSCFELTF
TSTSVAGKKMVVQVTINTGGDLGSSTGAHFDLQMPGGGVGIFNGCSSQWGAPNDGWGSR
YGGISSASDCSSLPALQAGCKWRFNWFKNADNPSMTYKEVTCPEITAKTGCSRK
(SEQ ID NO: 24)

[0103] In this amino acid sequence (SEQ ID NO: 24), amino acid residues 1 to 18 are signal peptides derived from NCE5, amino acid residues 19 to 24 are N-terminal sequences of LACE5, amino acid residues 25 to 36 are sequences derived from NCE5 or RCE I, and amino acid residue 37 and succeeding sequences are derived from the catalytic active domain of RCE I.

[0104] When the reaction is carried out in accordance with the above method, two types of sequences, i.e., the sequence as shown in SEQ ID NO: 24 and the sequence in which amino acid residue 34 is alanine, are obtained as amino acid sequences of the protein coded by the nucleotide sequence of the gene fragment H43. In the following procedure, a gene fragment used has a nucleotide sequence, which encodes the amino acid sequence as shown in SEQ ID NO: 24.

[Example 2] Construction of gene expressing RCE I variant H45 which lacks the cellulose-binding domain

[0105] pR1H4 obtained in Example 1 was used as a template to introduce amino acid substitution into the RCE I gene using synthetic DNA, RC-A121P, and the TaKaRa LA PCR in vitro Mutagenesis Kit (Takara Shuzo Co., Ltd.). Reaction conditions were in accordance with the manual attached to the kit. A sample was separated after the reaction by agarose gel electrophoresis to obtain an about 1-kbp gene fragment L9. This fragment was digested with the restriction enzyme *Bam*HI and ligated to the *Bam*HI site of plasmid pUC118 using the TaKaRa DNA Ligation Kit Ver. 1 to prepare plasmid pR1L9. Reaction conditions, such as those regarding enzymes, were in accordance with the manual attached to the kit. The obtained plasmid pR1L9 was subjected to sequencing reaction using the Cy5-Auto Read Sequencing Kit (Amersham Pharmacia), and the sequence was analyzed using the DNA Sequencer ALFred (Amersham Pharmacia). A synthetic DNA used as primer used in this reaction was H4-R1. As a result, it was confirmed that the sequence was in the expected form, i.e., one alanine was substituted with proline.

RC-A121P: GACTGCTGCAAGCCGTCGTGC (SEQ ID NO: 42);

5

H4-R1: GTTGCACATGTAGGAGTTGC (SEQ ID NO: 43)

[0106] Using this pR1L9 as a template, a gene encoding a portion of the secretion signal sequence in the RCE I gene was amplified. PCR was carried out using two synthetic DNAs, i.e., RC-451F and RC-451R, as primers, the TaKaRa Ex Taq Polymerase (Takara Shuzo Co., Ltd.), and attached buffers. The composition of the reaction solution was in accordance with the conditions specified in the attached manual. The temperature conditions for the Thermal Cycler (2400-R; Perkin Elmer) were 25 cycles of 94°C for 1 minute, 50°C for 2 minutes, and 72°C for 1 minute. A sample was separated after the reaction by agarose gel electrophoresis to obtain an about 100 bp gene fragment 45-X1.

[0107] Similarly, a region encoding the catalytic active domain (CAD) of the RCE I gene was amplified using the pR1L9 as a template. PCR was carried out using two synthetic DNAs, i.e., RC-452F and RC-452R, as primers, the TaKaRa Ex Taq Polymerase (Takara Shuzo Co., Ltd.), and attached buffers. The composition of the reaction solution was in accordance with the conditions specified in the attached manual. The temperature conditions for the Thermal Cycler (2400-R; Perkin Elmer) were 25 cycles of 94°C for 1 minute, 50°C for 2 minutes, and 72°C for 1 minute. A sample was separated after the reaction by agarose gel electrophoresis to obtain an about 630 bp gene fragment 45-X2.

20

RC-451F: GCGGATCCTGGGACAAGATG (SEQ ID NO: 25);

25

RC-451R: GCCTGCAGAGCGGCGGAGGCCATC (SEQ ID NO: 26);

RC-452F: GCCTGCAGGGAAAGTACAGCGCTGT (SEQ ID NO: 27);

30

RC-452R: GCGGATCCTGCGTTTACTTGC (SEQ ID NO: 28)

[0108] PCR fragments 45-X1 and 45-X2 were digested with the restriction enzyme *Pst*I, ligated together using the TaKaRa DNA Ligation Kit Ver. 1, and then digested with the restriction enzyme *Bam*HI to obtain a DNA fragment 45-X3. PCR was carried out using 45-X3 as a template, two synthetic DNAs, i.e., RC-451F and RC-452R, as primers, the TaKaRa Ex Taq Polymerase (Takara Shuzo Co., Ltd.), and attached buffers. The composition of the reaction solution was in accordance with the conditions specified in the attached manual. The temperature conditions for the Thermal Cycler (2400-R; Perkin Elmer) were 25 cycles of 94°C for 1 minute, 50°C for 2 minutes, and 72°C for 1 minute. A sample was separated after the reaction by agarose gel electrophoresis to obtain an about 700 bp gene fragment H45. This fragment was digested with the restriction enzyme *Bam*HI and ligated to the *Bam*HI site of plasmid pUC118 using the TaKaRa DNA Ligation Kit Ver. 1 to prepare plasmid pR1H45. Reaction conditions were in accordance with the manual attached to the kit. The amino acid sequence of the RCE I variant H45 deduced from the nucleotide sequence is shown below.

45

H45: MKFTTIASSALLALALGTEMASAALQGKYSVSGGASGNGVTTRYWDCKPSCSW

50

PGKANVSSPVKSCNKDGVLTALSDSNAQSGCNGGNSYMCNDNQPWAVNDNLAYGFAAA

AISGGGESRWCCSCFELTFTSTSVAGKKMVVQVTNTGGDLGSSTGAHFDLQMPGGGVGIF

55

NGCSSQWGAPNDGWGSGRYGGISSASDCSSLPSALQAGCKWRFNWFKNADNPSMTYKEV
TCPKEITAKTGCSRK (SEQ ID NO: 29)

[0109] In this amino acid sequence (SEQ ID NO: 29), amino acid residues 1 to 23 are signal peptides derived from

RCE1, amino acid residue 24 is an N-terminal sequence of RCE I, amino acid residues 25 and 26 are sequences introduced by a primer used, and amino acid residue 27 and succeeding sequences are derived from the catalytic active domain of RCE I.

- 5 [Example 3] Expression of codon optimized endoglucanase RCE I gene and its variants RCE I-H43 and H45, which lack the celluloses-binding domains, in *Humicola insolens*

[0110] Plasmid pJD01 (see Example D1 (2) (b) in WO 00/24879) was digested with BamHI and dephosphorylated using *Escherichia coli*-derived alkaline phosphatase (Takara Shuzo Co., Ltd.) in accordance with the attached manual. Plasmids pR1H43 and pR1H45 obtained in Examples 1 and 2 respectively were also digested with BamHI to produce about 700 bp DNA fragments. The obtained fragments were ligated respectively with aforementioned plasmid pJD01 using the DNA Ligation Kit Ver. 1 to produce expression plasmids pJND-H43 and pJND-H45.

[0111] *Humicola insolens* MN 200-1 (FERM BP-5977) was transformed with pJND-H43, pJND-H45, or pJ14D01 comprising the codon optimized RCE I gene (plasmid comprising a codon optimized RCE I gene to express the Rhizopus-derived RCE I in *Humicola*, see Example D3 (3) in WO 00/24879). Specifically, *Humicola insolens* MN 200-1 (FERM BP-5977) was cultured in medium (S) (3.0% glucose, 2.0% yeast extract, 0.1% polypeptone, 0.03% calcium chloride, 0.03% magnesium chloride (pH 6.8)) at 37°C for 24 hours, and then the cells were collected by centrifugation at 3,000 rpm for 10 minutes. The collected cells were washed with 0.5M sucrose and suspended in 10 ml of enzyme solution for preparing protoplast (5 mg/ml Novozyme 234 (Novo), 5 mg/ml Cellulase Onozuka R-10 (Yakult), and 0.5M sucrose) filtered through a 0.45 µm filter. The suspension was shaken at 30°C for 60 to 90 minutes to render fungal threads to be protoplasted. This suspension was filtered and then centrifuged at 2,500 rpm for 10 minutes, and protoplast was collected and washed with a SUTC buffer (0.5M sucrose, 10 mM calcium chloride, and 10 mM Tris hydrochloride (pH 7.5)).

[0112] The thus prepared protoplast was suspended in 1 mL of SUTC buffer, and 10 µl of DNA (TE) solution was added to the suspension in amounts of 10 µg per each 100 µl of the suspension. The mixture was allowed to stand in ice for 5 minutes. Subsequently, 400 µl of PEG solution (60% PEG 4000, 10 mM calcium chloride, and 10 mM Tris hydrochloride (pH 7.5)) was added and the mixture was allowed to stand in ice for 20 minutes. Thereafter, 10 ml of SUTC buffer was added, and centrifugation was carried out at 2,500 rpm for 10 minutes. The collected protoplast was suspended in 1 ml of SUTC buffer, centrifuged at 4,000 rpm for 5 minutes, and finally suspended in 100 µl of SUTC buffer.

[0113] The protoplast treated as above was superposed on a 200 µg/ml hygromycin B-containing YMG medium (1% glucose, 0.4% yeast extract, 0.2% malt extract, 1% agar (pH 6.8)) together with YMG soft agar. Culture was incubated at 37°C for 5 days. Thereafter, the generated colony was determined to be a transformant.

[0114] The obtained transformant was cultured in medium (N) (5.0% Avicel, 2.0% yeast extract, 0.1% polypeptone, 0.03% calcium chloride, 0.03% magnesium chloride, pH 6.8) at 37°C. A culture supernatant from which solid matter has been removed by centrifugation was determined to be an enzyme sample.

[Example 4] Isolation and purification of RCE I variant from *Humicola insolens* transformant

[0115] *Humicola insolens* transformants were inoculated to medium (N) (5.0% Avicel, 2.0% yeast extract, 0.1% polypeptone, 0.03% calcium chloride, 0.03% magnesium chloride, (pH 6.8)) and subjected to shake culture at 37°C. Transformants into which plasmids pJND-H43 and pJND-H45 had been introduced were cultured for 5 to 6 days. Regarding transformants into which plasmid pJ14D01 had been introduced, a sample was cultured for 4 days for obtaining an RCE I enzyme, which was not degraded in its linker domain, and which sustained a cellulose-binding domain. In contrast, a sample was cultured longer than 4 days, i.e., for 5 to 6 days, for obtaining the RCE I enzyme, which was degraded in its linker domain and lacked the cellulose-binding domain. Each of the resulting culture solutions was centrifuged at 7,000 rpm for 20 minutes to remove cells, and the culture supernatant was determined to be a crude cellulase preparation.

[0116] An ammonium sulfate solution at a final concentration of 1.5 M was prepared from 100 ml of this crude cellulase preparation and then applied at a flow rate of 10.0 ml/min to Macro-Prep Methyl HIC Support hydrophobic chromatography (270 ml in gel volume, BioRad Laboratories) which had been previously equilibrated with 1.5 M ammonium sulfate solution. It was then fractionated by eluting at a flow rate of 10.0 ml/min in a stepwise elution method in which the concentration of ammonium sulfate was decreased by 0.3 M each from 1.5 M. Fractions found to have strong activity of removing fuzz from lyocell were: a fraction obtained at an ammonium sulfate concentration of 1.2 M regarding a culture solution of the transformant into which plasmid pJND-H43 had been introduced (hereinafter referred to as an "H43 culture solution"); a fraction obtained at an ammonium sulfate concentration of 0.9 M regarding a culture solution of the transformant into which plasmid pJND-H45 had been introduced (hereinafter referred to as an "H45 culture solution"); a fraction obtained at an ammonium sulfate concentration of 0.6 M regarding a culture solution of the trans-

formant into which plasmid pJ14D01 had been introduced and cultured for 4 days (hereinafter referred to as an "H4 culture solution"); and a fraction obtained at an ammonium sulfate concentration of 0.9 M regarding a culture solution of the transformant into which plasmid pJ14D01 had been introduced and for which a culture period had been extended to accelerate its degradation (hereinafter referred to as an "H4 degradation product"). Therefore, 100 ml each of these fractions was fractionated.

[0117] An ammonium sulfate solution at a final concentration of 1.5 M was prepared from 100 ml of the obtained active fractions and then applied again at a flow rate of 10.0 ml/min to Macro-Prep Methyl HIC Support hydrophobic chromatography (270 ml in gel volume, BioRad Laboratories) which had been previously equilibrated with 1.5 M ammonium sulfate solution. It was then fractionated by eluting at a flow rate of 10.0 ml/min in a stepwise elution method in which the concentration of ammonium sulfate in deionized water was decreased by 0.15 M each from 1.5 M. Fractions found to have activities of removing fuzz from lycell were: a fraction obtained at an ammonium sulfate concentration of 1.35 M regarding the H43 culture solution; a fraction obtained at an ammonium sulfate concentration of 1.05 M regarding the H45 culture solution; a fraction obtained at an ammonium sulfate concentration of 0.75 M regarding the H4 culture solution; and a fraction obtained at an ammonium sulfate concentration of 1.05 M regarding the H4 degradation product. Therefore, 40 ml each of these fractions was fractionated.

[0118] An ammonium sulfate solution at a final concentration of 1.5 M was prepared from 40 ml of the obtained active fractions and then applied at a flow rate of 4.0 ml/min to Macro-Prep Methyl HIC Support hydrophobic chromatography (25 ml in gel volume, BioRad Laboratories) which had been previously equilibrated with 1.5 M ammonium sulfate solution. It was then fractionated by eluting at a flow rate of 4.0 ml/min in deionized water. Among these fractions, 8 ml of fraction, which was found to have strong activity of removing fuzz from lycell, was fractionated.

[0119] Acetate buffer (150 ml, 50 mM, pH 4.0) was prepared by diluting the obtained active fractions, and the resulting buffer was then applied at a flow rate of 2 ml/min to MonoS 10-/10-HR column (Amersham Pharmacia), which had been previously equilibrated with 50 mM acetate buffer (pH 4.0). It was then fractionated by eluting at a flow rate of 2 ml/min in a stepwise elution method in which the concentration of NaCl in 50 mM acetate buffer (pH 4.0) was increased by 0.1 M each to 1M NaCl in 50 mM acetate buffer (pH 5.0). Fractions that were obtained at a NaCl concentration of about 0.2 to 0.3M were found to have activity of removing fuzz from lycell. Therefore, 6 ml of fraction found to have the strongest activity was fractionated. These fractions showed in SDS-PAGE a single band of about 25 KDa regarding proteins purified from the H43 culture solution, the H45 culture solution, and the H4 degradation product and a single band of about 40 KDa regarding the protein purified from the H4 culture solution.

[0120] SDS-PAGE was carried out using the system of Tefco in which an electrophoresis tank (No. 03-101), a source (Model: 3540), 10% gel (01-015), and a buffer kit for SDS-PAGE (06-0301) were used. The condition for electrophoresis was 18 mA/10 min and then 20 mA/90 min. In protein detection after the electrophoresis, silver staining was carried out using a 2D-silver staining reagent IT "DAIICHI" (Daiichi Pure Chemicals Co., Ltd.) for electrophoresis. A standard protein used as a marker was the SDS-PAGE molecular weight standard protein, Low Range (161-0304, BioRad).

[0121] The activity of removing fuzz from lycell was measured in accordance with the following method.

[0122] Color knitted fabric of lycell (Toyoshima Japan) was fuzzed in a large washer together with a surfactant and rubber balls. Thereafter, the fuzzy knitted fabric of lycell (Toyoshima Japan, 9 cm x 10 cm, about 2 g in weight) was cylindrically sewn and subjected to fuzz removal treatment with various enzymes under the conditions as set forth below. The protein concentrations required to completely remove fuzz existing in the cylindrical fabric by this treatment were calculated.

[0123] The protein concentrations of various endoglucanases were calculated from the peak area at UV 280 nm of respective endoglucanase eluted with a linear gradient from 0% to 80% of acetonitrile concentration in 0.05% TFA (trifluoroacetic acid) at a flow rate of 1.0 ml/min in HPLC analysis using TSK gel TMS-250 column (4.6 mm LD. x 7.5 cm, TOSOH Japan). The standard used was the purified NCE4, which was analyzed in HPLC under the same conditions, the protein concentration of which had been preliminarily measured by the protein assay kit (BioRad Laboratories). The standard used to measure the protein concentration for the protein assay kit was albumin standard (Bovine serum albumin, fraction V, PIERCE). The purified NCE4 (coded by nucleotides 118 to 1088 in SEQ ID NO: 18) was isolated and purified from a culture solution of *Humicola insolens* according to the method described in WO 98/03640.

Test machine: Launder Meter L-12 (Daiei Kagaku Seiki MFG, Japan)

Temperature: 55°C

Time: 60 minutes

Reaction volume: 40 ml

Reaction pH: pH 5 (10 mM acetate buffer)

pH 6 (10 mM acetate buffer)

[0124] The treating liquid contained 4 rubber balls (about 16 g each) together with the endoglucanase solution.

[Example 5] Identification of N-terminal amino acid sequences of RCE I variant protein isolated and purified from *Humicola insolens* transformant

5 [0125] In order to determine the N-terminal amino acid sequences of the purified proteins obtained in Example 4, each of the fractions was subjected to SDS-PAGEmini (Tefco), electroblotted on a PVDF membrane, stained with Coomassie Brilliant Blue R-250 (Nacalai Tesque, Inc.), decolorized, washed with water, and then air dried. A portion on which the object protein had been blotted was cleaved out therefrom and subjected to a Protein Sequencer (Model 492, PE Applied Biosystems) to analyze the N-terminal amino acid sequence. The amino acid sequences were read from enzymes that were purified from the H45 culture solution, the H4 culture solution, and the H4 degradation product 10 without any problem. Regarding the enzyme purified from the H43 culture solution, however, no signal was generated by Edman degradation and the N-terminal amino acid was found to be modified and protected. Thus, this enzyme was immersed in a solution of 0.5% polyvinyl pyrrolidone (molecular weight: 40,000, Sigma)/100 mM acetic acid at 37°C for 30 minutes, and, after a protein unbound portion on the membrane was blocked, this enzyme was treated with pfu Pyroglutamate Aminopeptidase (Takara Shuzo Co., Ltd.) at 50°C for 5 hours to remove the modified N-terminal residue. 15 Thus, sequencing was carried out once more. The obtained sequences were as shown below.

The N-terminal amino acid sequence of RCE I-H43:

20 [0126]

Gln-Ser-Gly-Ser-Gly-Arg-Thr (7 residues) (SEQ ID NO: 30).

25 The N-terminal amino acid sequence of RCE I-H45:

[0127]

30 **Lys-Tyr-Ser-Ala-Val-Ser-Gly (7 residues) (SEQ ID NO: 31);**

and

35 **Ala-Val-Ser-Gly-Gly-Ala-Ser (7 residues) (SEQ ID NO: 32).**

The N-terminal amino acid sequence of RCE I-H4(25KDa):

40 [0128]

Ser-Ala-Val-Ser-Gly-Gly-Ala (7 residues) (SEQ ID NO: 33);

45 and

50 **Gly-Gly-Ala-Ser-Gly-Asn-Gly (7 residues) (SEQ ID NO: 34).**

The N-terminal amino acid sequence of RCE I-H4 (40KDa):

[0129]

55 **Ala-Glu-(Cys)-Ser-Lys-Leu-Tyr (7 residues) (SEQ ID NO: 35).**

[0130] As a result of identification of N-terminal amino acid sequences, it was found that only the enzyme purified from the H4 culture solution (hereinafter referred to as "RCE I-H4 (40 KDa)") had a cellulose-binding domain (CBD) while any of the remaining enzyme purified from the H43 culture solution (hereinafter referred to as "RCE I-H43 (25 KDa)"), the enzyme purified from the H45 culture solution (hereinafter referred to as "RCE I-H45 (25 KDa)"), or the enzyme purified from the H4 degradation product (hereinafter referred to as "RCE I-H4 (25 KDa)") had no cellulose-binding domain (CBD) but had only the catalytic active domains (CAD).

[Example 6] Comparison of specific activity of removing fuzz from cotton fabric between RCE I, which lacks the cellulose-binding domain, and RCE L which has the cellulose-binding domain

[0131] Using the endoglucanase which was homogenously purified in Example 5, a knitted cotton fabric (a 6 cm x 8 cm fabric from cotton smooth knit No. 3900, Nitto Boseki Co., Ltd., dyed in brown by reactive dye at Tsuyatomo Senko), which has been fuzzed in a large washer, was subjected to fuzz removal under the following conditions. The amount of fuzz remaining unremoved was visually evaluated, and the amount of purified enzyme to be added so that the amount of remaining fuzz would reach 50% was determined. The amount of protein was determined using the BCA Protein Assay Reagent (PIERCE) in accordance with the conditions described in the attached manual. The estimated molecular weight of the 40 KDa purified RCE I protein (RCE I-H4 (40 KDa)) is about 1.5 times higher than those of the 25 KDa purified RCE I proteins (RCE I-H43 (25 KDa)), RCE I-H45 (25 KDa)), and RCE I-H4 (25 KDa)). Accordingly, even though the quantified amounts of proteins are the same, the number of enzyme molecules in the 40 KDa purified RCE I protein is only about two-thirds of that in the 25 KDa purified RCE I protein in terms of the number of enzyme molecules contained in the protein.

Test machine: Launder Meter L-20 (Daiei Kagaku Seiki MFG, Japan)

Temperature: 40°C or 55°C

Time: 120 minutes

Reaction volume: 40 ml

Reaction pH: Reacted at pH 7 (1 mM phosphate buffer, prepared using deionized water).

[0132] The treating liquid contained 4 rubber balls (about 16 g each) together with the enzyme solution.

[0133] The results are as shown in Table 2 below.

Table 2

	The amount of enzyme added 40°C	The amount of enzyme added 55°C
Purified RCE I-H4 (40 KDa)	390 µg or more	390 µg or more
Purified RCE I-H4 (25 KDa)	18 µg	53 µg

[0134] As is apparent from the results shown in Table 2, the 25 KDa protein, which lacks the cellulose-binding domain (CBD), exhibits much higher activity of removing fuzz from cotton fabrics than the 40 KDa protein, which has the cellulose-binding domain (CBD), even though both proteins are originated from the same *Zygomycetes*-derived endoglucanases RCE I.

[Example 7] Comparison of specific activity of removing fuzz from lyocell fabric between RCE I, which lacks the cellulose-binding domain, and RCE L, which has the cellulose-binding domain

[0135] Using the endoglucanase which was homogenously purified in Example 5, a lyocell fabric (6 cm x 8 cm, Toyoshima Japan) was subjected to fuzz removal under conditions improved from the method described in Example 4. The amount of fuzz remaining unremoved was visually evaluated, and the amount of purified enzyme required to completely remove fuzz was determined. The amount of protein was determined using the BCA Protein Assay Reagent (PIERCE) in accordance with the conditions described in the attached manual. The estimated molecular weight of the 40 KDa purified RCE I protein (RCE I-H4 (40 KDa)) is about 1.5 times higher than those of the 25 KDa purified RCE I proteins (RCE I-H43 (25 KDa)), RCE I-H45 (25 KDa)), and RCE I-H4 (25 KDa)). Accordingly, even though the quantified amounts of proteins are the same, the number of enzyme molecules in the 40 KDa purified RCE I protein is only about two-thirds of that in the 25 KDa purified RCE I protein in terms of the number of enzyme molecules contained in the protein. Test machine: Launder Meter L-20 (Daiei Kagaku Seiki MFG, Japan)

Temperature: 40°C

Time: 90 minutes

Reaction volume: 50 ml

Reaction pH: Reacted at pH 6 (10 mM acetate buffer, prepared using deionized water).

[0136] The treating liquid contained 4 rubber balls (about 16 g each) together with the enzyme solution.

[0137] The results are as shown in Table 3 below.

Table 3

	The amount of enzyme added
Purified RCE I-H4 (40 KDa)	32 µg
Purified RCE I-H4 (25 KDa)	11 µg

[0138] As is apparent from the results shown in Table 3, RCE I, which is Zygomycetes-derived endoglucanase, exhibits a higher specific activity of removing fuzz from lyocell fabrics as the 25 KDa protein, which lacks the cellulose-binding domain (CBD), than as the 40 KDa protein, which has the cellulose-binding domain (CBD).

[Example 8] Comparison of specific activity of removing fuzz from cotton fabric between RCE I, which lacks the cellulose-binding domain, and RCE I, which has the cellulose-binding domain, under alkaline, low-temperature, and surfactant-present conditions

[0139] Using the endoglucanase which was homogenously purified in Example 5, a knitted cotton fabric (a 6 cm x 8 cm fabric from cotton smooth knit No. 3900, Nitto Boseki Co., Ltd., dyed in brown by reactive dye at Tsuyatomo Senko), which had been fuzzed in a large washer, was subjected to fuzz removal under the following conditions. The amount of fuzz remaining unremoved was visually evaluated, and the amount of purified enzyme to be added so that the amount of remaining fuzz would reach 50% was determined. The amount of protein was determined using the BCA Protein Assay Reagent (PIERCE) in accordance with the conditions described in the attached manual. The estimated molecular weight of the 40 KDa purified RCE I protein (RCE I-H4 (40 KDa)) is about 1.5 times higher than those of the 25 KDa purified RCE I proteins (RCE I-H43 (25 KDa)), RCE I-H45 (25 KDa)), and RCE I-H4 (25 KDa)). Accordingly, even though the quantified amounts of proteins are the same, the number of enzyme molecules in the 40 KDa purified RCE I protein is only about two-thirds of that in the 25 KDa purified RCE I protein in terms of the number of enzyme molecules contained in the protein.

Test machine: Launder Meter L-20 (Daiei Kagaku Seiki MFG, Japan)

Temperature: 30°C

Time: 120 minutes

Reaction volume: 40 ml

Reaction pH: Reacted at pH 10 (5 mM sodium carbonate buffer, prepared using deionized water).

[0140] The treating liquid contained a nonionic surfactant Persoft NK-100 (NOF Corp.) at a final concentration of 100 ppm together with the enzyme solution and 4 rubber balls (about 16 g each).

[0141] The results are as shown in Table 4 below.

Table 4

	The amount of enzyme added
Purified RCE I-H4 (40 KDa)	390 µg or more
Purified RCE I-H4 (25 KDa)	52 µg

[0142] As is apparent from the results shown in Table 4, given the low-temperature, alkaline, and surfactant-present conditions under which detergents are actually used, the 25 KDa protein, which lacks the cellulose-binding domain (CBD), exhibits much higher activity of removing fuzz from cotton fabrics than the 40 KDa protein, which has the cellulose-binding domain (CBD), even though both proteins are originated from the same Zygomycetes-derived endoglucanases RCE I.

[Example 9] Comparison of specific activity of removing fuzz from cotton fabric among various purified RCE I proteins, which lack the cellulose-binding domains

[0143] Using the endoglucanase which was homogenously purified in Example 5, a knitted cotton fabric (a 6 cm x 8 cm fabric from cotton smooth knit No. 3900, Nitto Boseki Co., Ltd., dyed in brown by reactive dye at Tsuyatomo Senko), which had been fuzzed in a large washer, was subjected to fuzz removal under the following conditions. The amount of fuzz remaining unremoved was visually evaluated, and the amount of purified enzyme to be added so that the amount of remaining fuzz would reach 50% was determined. The amount of protein was determined using the BCA Protein Assay Reagent (PIERCE) in accordance with the conditions described in the attached manual. The estimated

molecular weight of the 40 KDa purified RCE I protein (RCE I-H4 (40 KDa)) is about 1.5 times higher than those of the 25 KDa purified RCE I proteins (RCE I-H43 (25 KDa)), RCE I-H45 (25 KDa)), and RCE I-H4 (25 KDa)). Accordingly, even though the quantified amounts of proteins are the same, the number of enzyme molecules in the 40 KDa purified RCE I protein is only about two-thirds of that in the 25 KDa purified RCE I protein in terms of the number of enzyme molecules contained in the protein.

Test machine: Launch Meter L-20 (Daiei Kagaku Seiki MFG, Japan)

Temperature: 40°C

Time: 120 minutes

Reaction volume: 40 ml

Reaction pH: Reacted at pH 7 (1 mM phosphate buffer, prepared using deionized water).

[0144] The treating liquid contained 4 rubber balls (about 16 g each) together with the enzyme solution.

[0145] The results are as shown in Table 5 below.

Table 5

	The amount of enzyme added
Purified RCE I-H4 (40 KDa)	390 µg or more
Purified RCE I-H43 (25 KDa)	26 µg
Purified RCE I-H45 (25 KDa)	18 µg
Purified RCE I-H4 (25 KDa)	18 µg

[0146] As is apparent from the results shown in Table 5, various RCE I proteins such as RCE I-H43 (25 KDa), RCE I-H45 (25 KDa), and RCE I-H4 (25 KDa), which lack the cellulose-binding domains, exhibit much higher activity of removing fuzz from cotton fabrics than the 40 KDa protein RCE I-H4 (40 KDa), which has the cellulose-binding domain (CBD). This indicates that the RCE I protein, which lacks the cellulose-binding domain (CBD), exhibits much higher activity of removing fuzz from cotton fabrics than the 40 KDa protein, which has the cellulose-binding domain (CBD), regardless the length of the linker domain remaining on the N-terminal side of the catalytic active domain. Also, whether or not it is an artificial or non-artificial deficiency, the RCE I protein, which lacks the cellulose-binding domain (CBD), exhibits much higher activity of removing fuzz from cotton fabrics than the 40 KDa protein RCE I-H4 (40 KDa), which has the cellulose-binding domain (CBD).

[Example 10] Comparison of specific activity of removing fuzz from lyocell fabric among various purified RCE I proteins, which lack the cellulose-binding domains

[0147] Using the endoglucanase which was homogenously purified in Example 5, a lyocell fabric (6 cm x 8 cm, Toyoshima Japan) was subjected to fuzz removal under conditions improved from the method described in Example 4. The amount of fuzz remaining unremoved was visually evaluated, and the amount of purified enzyme required to completely remove fuzz was determined. The amount of protein was determined using the BCA Protein Assay Reagent (PIERCE) in accordance with the conditions described in the attached manual. The estimated molecular weight of the 40 KDa purified RCE I protein (RCE I-H4 (40 KDa)) is about 1.5 times higher than those of the 25 KDa purified RCE I proteins (RCE I-H43 (25 KDa)), RCE I-H45 (25 KDa)), and RCE I-H4 (25 KDa)). Accordingly, even though the quantified amounts of proteins are the same, the number of enzyme molecules in the 40 KDa purified RCE I protein is only about two-thirds of that in the 25 KDa purified RCE I protein in terms of the number of enzyme molecules contained in the protein.

Test machine: Launder Meter L-20 (Daiei Kagaku Seiki MFG, Japan)

Temperature: 40°C

Time: 90 minutes

Reaction volume: 40 ml

Reaction pH: Reacted at pH 6 (10 mM acetate buffer, prepared using deionized water).

[0148] The treating liquid contained 4 rubber balls (about 16 g each) together with the enzyme solution.

[0149] The results are as shown in Table 6 below.

Table 6

	The amount of enzyme added
Purified RCE I-H4 (40 KDa)	32 µg
Purified RCE I-H43 (25 KDa)	12 µg

Table 6 (continued)

	The amount of enzyme added
Purified RCE I-H45 (25 KDa)	11 µg
Purified RCE I-H4 (25 KDa)	11 µg

[0150] As is apparent from the results shown in Table 6, various RCE I proteins such as RCE I-H43 (25 KDa), RCE I-H-45 (25 KDa), and RCE I-H4 (25 KDa), which lack the cellulose-binding domains, exhibit higher activity of removing fuzz from lyocell fabrics than the 40 KDa protein RCE I-H4 (40 KDa), which has the cellulose-binding domain (CBD). This indicates that the RCE I protein, which lacks the cellulose-binding domain (CBD), exhibits higher activity of removing fuzz from lyocell fabrics than the 40 KDa protein, which has the cellulose-binding domain (CBD) regardless the length of the linker domain remaining on the N-terminal side of the catalytic active domain. Also, whether or not it is an artificial or non-artificial deficiency, the RCE I protein, which lacks the cellulose-binding domain (CBD), exhibits higher activity of removing fuzz from lyocell fabrics than the 40 KDa protein RCE I-H4 (40 KDa), which has the cellulose-binding domain (CBD).

[0151] All publications, patents, and patent applications cited herein are incorporated herein by reference in their entirety.

Industrial Applicability

[0152] When Zygomycetes-derived endoglucanase such as RCE I, RCE II, RCE III, MCE I, MCE II, or PCE I, which lacks the cellulose-binding domain, is allowed to act, the endoglucanase activity can be significantly enhanced in comparison with endoglucanase having a cellulose-binding domain. Accordingly, fabric treatment such as reduction of fuzzing, improvement in feel and appearance, color clarification, partial color change, and softening of cellulose-containing fabrics and improvement in deinking of waste paper and drainage of paper pulp can be effected with a smaller amount of enzymes. This can decrease necessary costs remarkably.

Free Text of Sequence Listings

[0153]

SEQ ID NO: 13: codon optimized sequence corresponding to RCE I protein (SEQ ID NO: 2)
 SEQ ID NO: 17: consensus amino acid sequence found in the cellulose-binding domain of family 45 endoglucanase
 SEQ ID NO: 20 to 23: primers
 SEQ ID NO: 24: recombinant protein
 SEQ ID NO: 25 to 28: primers
 SEQ ID NO: 29: recombinant protein
 SEQ ID NO: 30 to 35: N-terminal amino acid sequence of recombinant protein
 SEQ ID NO: 36 and 37: primers
 SEQ ID NO: 38: NCE5 amino acid sequence
 SEQ ID NO: 39: NCE5 cDNA sequence
 SEQ ID NO: 40 to 43: primers

SEQUENCE LISTING

5

<110> MEIJI SEIKA KAISHA, LTD.

10

<120> Endoglucanase Enzyme from Zygomycetes Having Deletion of
Cellulose Binding Domain

15

<130> PH-1443-PCT

20

<150> JP2000-354296

<151> 2000-11-21

25

<160> 43

30

<170> PatentIn Ver. 2.0

<210> 1

35

<211> 338

<212> PRT

<213> Rhizopus oryzae CP96001

40

<220>

45

<221> sig#peptide

<222> (-23)... (-1)

50

<220>

<221> mat#peptide

55

<222> (1)... (315)

<400> 1

5 Met Lys Phe Ile Thr Ile Ala Ser Ser Ala Leu Leu Ala Leu Ala Leu
 -20 -15 -10
 Gly Thr Glu Met Ala Ser Ala Ala Glu Cys Ser Lys Leu Tyr Gly Gln
 10 -5 1 5
 Cys Gly Gly Lys Asn Trp Asn Gly Pro Thr Cys Cys Glu Ser Gly Ser
 10 15 20 25
 15 Thr Cys Lys Val Ser Asn Asp Tyr Tyr Ser Gln Cys Leu Pro Ser Gly
 30 35 40
 20 Ser Ser Gly Asn Lys Ser Ser Glu Ser Ala His Lys Lys Thr Thr Thr
 45 50 55
 Ala Ala His Lys Lys Thr Thr Thr Ala Ala His Lys Lys Thr Thr Thr
 25 60 65 70
 Ala Pro Ala Lys Lys Thr Thr Thr Val Ala Lys Ala Ser Thr Pro Ser
 75 80 85
 30 Asn Ser Ser Ser Ser Ser Ser Gly Lys Tyr Ser Ala Val Ser Gly Gly
 90 95 100 105
 35 Ala Ser Gly Asn Gly Val Thr Thr Arg Tyr Trp Asp Cys Cys Lys Ala
 110 115 120
 Ser Cys Ser Trp Pro Gly Lys Ala Asn Val Ser Ser Pro Val Lys Ser
 40 125 130 135
 Cys Asn Lys Asp Gly Val Thr Ala Leu Ser Asp Ser Asn Ala Gln Ser
 140 145 150
 45 Gly Cys Asn Gly Gly Asn Ser Tyr Met Cys Asn Asp Asn Gln Pro Trp
 155 160 165
 50 Ala Val Asn Asp Asn Leu Ala Tyr Gly Phe Ala Ala Ala Ala Ile Ser
 170 175 180 185
 Gly Gly Gly Glu Ser Arg Trp Cys Cys Ser Cys Phe Glu Leu Thr Phe
 55 190 195 200

Thr Ser Thr Ser Val Ala Gly Lys Lys Met Val Val Gln Val Thr Asn
 205 210 215
 Thr Gly Gly Asp Leu Gly Ser Ser Thr Gly Ala His Phe Asp Leu Gln
 220 225 230
 Met Pro Gly Gly Gly Val Gly Ile Phe Asn Gly Cys Ser Ser Gln Trp
 235 240 245
 Gly Ala Pro Asn Asp Gly Trp Gly Ser Arg Tyr Gly Gly Ile Ser Ser
 250 255 260 265
 Ala Ser Asp Cys Ser Ser Leu Pro Ser Ala Leu Gln Ala Gly Cys Lys
 270 275 280
 Trp Arg Phe Asn Trp Phe Lys Asn Ala Asp Asn Pro Ser Met Thr Tyr
 285 290 295
 Lys Glu Val Thr Cys Pro Lys Glu Ile Thr Ala Lys Thr Gly Cys Ser
 300 305 310
 Arg Lys
 315

<210> 2
 <211> 1017
 <212> DNA
 <213> Rhizopus oryzae CP96001

<220>
 <221> sig#peptide
 <222> (1)... (69)

<220>
 <221> mat#peptide
 <222> (70)... (1017)

<400> 2

5	atg aag ttt att act att gcc tct tcc gct ctc ttg gct ctc gcc ctc	48
	Met Lys Phe Ile Thr Ile Ala Ser Ser Ala Leu Leu Ala Leu Ala Leu	
10	-20 -15 -10	
	ggt act gaa atg gcc tct gct gct gaa tgt agc aaa ttg tat ggt caa	96
	Gly Thr Glu Met Ala Ser Ala Ala Glu Cys Ser Lys Leu Tyr Gly Gln	
15	-5 1 5	
	tgt ggt ggt aag aac tgg aat ggc cct act tgt tgt gaa tct gga tcc	144
20	Cys Gly Gly Lys Asn Trp Asn Gly Pro Thr Cys Cys Glu Ser Gly Ser	
	10 15 20 25	
	acc tgt aaa gta agc aac gat tac tac tct caa tgt ctt ccc tct gga	192
25	Thr Cys Lys Val Ser Asn Asp Tyr Tyr Ser Gln Cys Leu Pro Ser Gly	
	30 35 40	
	agc agt ggc aat aaa tct tct gaa agt gct cac aag aag act acc act	240
30	Ser Ser Gly Asn Lys Ser Ser Glu Ser Ala His Lys Lys Thr Thr Thr	
	45 50 55	
	gct gct cac aag aag act act acc gct gct cat aaa aag act acc act	288
35	Ala Ala His Lys Lys Thr Thr Thr Ala Ala His Lys Lys Thr Thr Thr	
	60 65 70	
40	gct cct gct aag aag act aca act gtt gcc aaa gct tcc acc cct tct	336
	Ala Pro Ala Lys Lys Thr Thr Thr Val Ala Lys Ala Ser Thr Pro Ser	
	75 80 85	
45	aac tct agc tct agc tcc agc ggc aaa tat tcc gct gtc tct ggt ggt	384
	Asn Ser Ser Ser Ser Ser Ser Gly Lys Tyr Ser Ala Val Ser Gly Gly	
	90 95 100 105	
50	gcc tct ggt aac ggt gtc act act cgt tat tgg gat tgc tgt aag gcc	432
	Ala Ser Gly Asn Gly Val Thr Thr Arg Tyr Trp Asp Cys Cys Lys Ala	
55	110 115 120	

tcc tgt agc tgg ccc ggt aag gcc aat gtc agt tct cct gtc aag tcc 480
 Ser Cys Ser Trp Pro Gly Lys Ala Asn Val Ser Ser Pro Val Lys Ser
 5 125 130 135
 tgt aac aaa gat ggt gtc act gcc ctt agt gac agc aat gcc caa agt 528
 10 Cys Asn Lys Asp Gly Val Thr Ala Leu Ser Asp Ser Asn Ala Gln Ser
 140 145 150
 ggc tgt aac ggt ggt aac agt tac atg tgt aac gac aac caa cct tgg 576
 15 Gly Cys Asn Gly Gly Asn Ser Tyr Met Cys Asn Asp Asn Gln Pro Trp
 155 160 165
 gct gta aac gac aac ctt gcc tat ggt ttc gct gct gct gcc atc agt 624
 20 Ala Val Asn Asp Asn Leu Ala Tyr Gly Phe Ala Ala Ala Ala Ile Ser
 170 175 180 185
 ggt ggt ggt gaa tct cgc tgg tgc tgt tct tgt ttc gaa ctt act ttc 672
 25 Gly Gly Gly Glu Ser Arg Trp Cys Cys Ser Cys Phe Glu Leu Thr Phe
 190 195 200
 act tct acc tct gtt gct ggt aag aag atg gtt gtc caa gtc act aac 720
 30 Thr Ser Thr Ser Val Ala Gly Lys Lys Met Val Val Gln Val Thr Asn
 205 210 215
 act ggt ggt gat ctt ggc tcc tct act ggt gct cac ttt gac ttg caa 768
 35 Thr Gly Gly Asp Leu Gly Ser Ser Thr Gly Ala His Phe Asp Leu Gln
 220 225 230
 atg ccc ggt ggt ggt gtt ggt att ttc aat ggt tgt tcc agc caa tgg 816
 40 Met Pro Gly Gly Gly Val Gly Ile Phe Asn Gly Cys Ser Ser Gln Trp
 235 240 245
 ggt gct ccc aat gac ggt tgg ggc tca aga tac ggt ggt att tct tct 864
 50 Gly Ala Pro Asn Asp Gly Trp Gly Ser Arg Tyr Gly Gly Ile Ser Ser
 250 255 260 265
 gca tct gac tgc tct agt ctt cct tcc gca ctc caa gct ggt tgt aaa 912
 55 Ala Ser Asp Cys Ser Ser Leu Pro Ser Ala Leu Gln Ala Gly Cys Lys

270 275 280
 5 tgg aga ttc aac tgg ttc aag aac gct gat aac cca agc atg act tac 960
 Trp Arg Phe Asn Trp Phe Lys Asn Ala Asp Asn Pro Ser Met Thr Tyr
 285 290 295
 10 aag gaa gtt acc tgt cct aag gaa atc acc gcc aag aca ggt tgt tca 1008
 Lys Glu Val Thr Cys Pro Lys Glu Ile Thr Ala Lys Thr Gly Cys Ser
 300 305 310
 15 aga aaa taa 1017
 Arg Lys
 315
 20
 <210> 3
 25 <211> 366
 <212> PRT
 <213> Rhizopus oryzae CP96001
 30
 <220>
 <221> sig#peptide
 35 <222> (-23)... (-1)
 <220>
 40 <221> mat#peptide
 <222> (1)... (343)
 45
 <400> 3
 50 Met Lys Phe Ile Thr Ile Thr Ser Ser Ala Leu Leu Ala Leu Ala Leu
 -20 -15 -10
 Gly Thr Glu Met Ala Ser Ala Ala Lys Cys Ser Lys Leu Tyr Gly Gln
 55 -5 1 5

	Cys Gly Gly Lys Asp Trp Asn Gly Pro Thr Cys Cys Glu Ser Gly Ser	
5	10 15 20 25	
	Thr Cys Lys Val Ser Asn Asp Tyr Tyr Ser Gln Cys Leu Ala Pro Glu	
	30 35 40	
10	Ser Asn Gly Asn Lys Ser Ser Glu Cys Ser Lys Leu Tyr Gly Gln Cys	
	45 50 55	
	Gly Gly Lys Asp Trp Asn Gly Pro Thr Cys Cys Glu Ser Gly Ser Thr	
15	60 65 70	
	Cys Lys Val Ser Asn Asp Tyr Tyr Ser Gln Cys Leu Ala Pro Glu Ser	
20	75 80 85	
	Asn Gly Asn Lys Thr Ser Glu Ser Ala His Lys Thr Thr Thr Thr Thr	
	90 95 100 105	
25	Ala Pro Ala Lys Glu Ile Thr Thr Thr Ala Lys Ala Ser Asn Ser Ser	
	110 115 120	
	Asn Ser Ser Gly Lys Tyr Ser Ile Val Ser Gly Gly Ala Ser Gly Asn	
30	125 130 135	
	Gly Val Thr Thr Arg Tyr Trp Asp Cys Cys Lys Ala Ser Cys Ser Trp	
35	140 145 150	
	Pro Gly Lys Ala Asn Val Ser Ser Pro Val Lys Ser Cys Asn Lys Asp	
	155 160 165	
40	Gly Val Thr Ala Leu Ser Asp Ser Asn Val Gln Ser Gly Cys Asn Gly	
	170 175 180 185	
	Gly Asn Ser Tyr Met Cys Asn Asp Asn Gln Pro Trp Ala Val Asn Asp	
45	190 195 200	
	Asn Leu Ala Tyr Gly Phe Ala Ala Ala Ala Ile Ser Gly Gly Gly Glu	
50	205 210 215	
	Ser Arg Trp Cys Cys Ser Cys Phe Glu Leu Thr Phe Thr Ser Thr Ser	
	220 225 230	
55	Val Ala Gly Lys Lys Met Val Ile Gln Val Thr Asn Thr Gly Gly Asp	

235 240 245
 5 Leu Gly Ser Ser Thr Gly Ala His Phe Asp Leu Gln Met Pro Gly Gly
 250 255 260 265
 Gly Val Gly Ile Phe Asn Gly Cys Ser Lys Gln Trp Gly Ala Pro Asn
 10 270 275 280
 Asp Gly Trp Gly Ser Arg Tyr Gly Gly Ile Ser Ser Ala Ser Asp Cys
 285 290 295
 15 Ser Ser Leu Pro Ser Ala Leu Gln Ala Gly Cys Lys Trp Arg Phe Asn
 300 305 310
 20 Trp Phe Lys Asn Ala Asp Asn Pro Ser Met Thr Tyr Lys Glu Val Thr
 315 320 325
 Cys Pro Lys Glu Ile Thr Ala Lys Thr Gly Cys Ser Arg Lys
 25 330 335 340

 30 <210> 4
 <211> 1101
 <212> DNA
 35 <213> Rhizopus oryzae CP96001

 40 <220>
 <221> sig#peptide
 <222> (1)... (69)

 45 <220>
 <221> mat#peptide
 <222> (70)... (1101)
 50

 55 <400> 4
 atg aag ttt att act att acc tct tcc gct ctc ttg gct ctc gcc ctt 48

	Met Lys Phe Ile Thr Ile Thr Ser Ser Ala Leu Leu Ala Leu Ala Leu			
	-20	-15	-10	
5	ggc act gaa atg gcc tct gct gct aaa tgt agc aag ctg tat ggt caa	96		
	Gly Thr Glu Met Ala Ser Ala Ala Lys Cys Ser Lys Leu Tyr Gly Gln			
10	-5	1	5	
	tgt ggt ggt aag gac tgg aat ggc cct act tgt tgc gaa tct gga tcc	144		
	Cys Gly Gly Lys Asp Trp Asn Gly Pro Thr Cys Cys Glu Ser Gly Ser			
15	10	15	20	25
	acc tgt aaa gta agc aac gat tac tac tct caa tgt ctt gcc cct gaa	192		
20	Thr Cys Lys Val Ser Asn Asp Tyr Tyr Ser Gln Cys Leu Ala Pro Glu			
	30	35	40	
	agc aac ggc aat aag tct tct gaa tgt agc aag ttg tat ggt caa tgt	240		
25	Ser Asn Gly Asn Lys Ser Ser Glu Cys Ser Lys Leu Tyr Gly Gln Cys			
	45	50	55	
	ggc ggt aag gac tgg aat ggc cct act tgt tgc gaa tct gga tcc acc	288		
30	Gly Gly Lys Asp Trp Asn Gly Pro Thr Cys Cys Glu Ser Gly Ser Thr			
	60	65	70	
35	tgt aaa gla agc aac gat tac tac tct caa tgt ctt gcc cct gaa agc	336		
	Cys Lys Val Ser Asn Asp Tyr Tyr Ser Gln Cys Leu Ala Pro Glu Ser			
	75	80	85	
40	aat ggc aat aaa act tct gaa agc gct cat aaa acg act act acc act	384		
	Asn Gly Asn Lys Thr Ser Glu Ser Ala His Lys Thr Thr Thr Thr Thr			
	90	95	100	105
45	gct ccc gct aag gaa att aca act act gcc aaa gct tca aac tct tct	432		
	Ala Pro Ala Lys Glu Ile Thr Thr Thr Ala Lys Ala Ser Asn Ser Ser			
	110	115	120	
50	aac tct agc ggc aaa tac tcc att gtc tct ggt ggt gcc tct ggt aac	480		
	Asn Ser Ser Gly Lys Tyr Ser Ile Val Ser Gly Gly Ala Ser Gly Asn			
55	125	130	135	

	ggc gtc act act cgt tat tgg gat tgc tgt aag gcc tcc tgt agc tgg	528
5	Gly Val Thr Thr Arg Tyr Trp Asp Cys Cys Lys Ala Ser Cys Ser Trp	
	140 145 150	
	ccc ggt aag gcc aat gtc agt tct cct gtc aag tcc tgt aac aaa gat	576
10	Pro Gly Lys Ala Asn Val Ser Ser Pro Val Lys Ser Cys Asn Lys Asp	
	155 160 165	
	ggc gtc act gcc ctt agt gac agc aat gtc caa agt ggc tgt aac ggt	624
15	Gly Val Thr Ala Leu Ser Asp Ser Asn Val Gln Ser Gly Cys Asn Gly	
	170 175 180 185	
	ggc aac agt tac atg tgt aac gac aac cag cct tgg gct gta aac gat	672
20	Gly Asn Ser Tyr Met Cys Asn Asp Asn Gln Pro Trp Ala Val Asn Asp	
	190 195 200	
	aat ctt gcc tat ggt ttc gct gct gct gcc atc agt ggt ggt ggt gaa	720
25	Asn Leu Ala Tyr Gly Phe Ala Ala Ala Ala Ile Ser Gly Gly Gly Glu	
	205 210 215	
	tct cgc tgg tgc tgt tct tgt ttc gaa ctt act ttc act tct acc tct	768
30	Ser Arg Trp Cys Cys Ser Cys Phe Glu Leu Thr Phe Thr Ser Thr Ser	
	220 225 230	
	ggt gct ggt aag aag atg gtt atc caa gtc act aac act ggt ggt gat	816
35	Val Ala Gly Lys Lys Met Val Ile Gln Val Thr Asn Thr Gly Gly Asp	
	235 240 245	
	ctt ggc tcc tct act ggt gct cac ttt gac ttg caa atg ccc ggt ggt	864
40	Leu Gly Ser Ser Thr Gly Ala His Phe Asp Leu Gln Met Pro Gly Gly	
	250 255 260 265	
	ggc gtt ggt att ttc aat ggt tgc tcc aag caa tgg ggt gct ccc aat	912
45	Gly Val Gly Ile Phe Asn Gly Cys Ser Lys Gln Trp Gly Ala Pro Asn	
	270 275 280	
	gac ggt tgg ggc tgc aga tac ggt ggt att tct tct gca tct gac tgc	960
50	Asp Gly Trp Gly Ser Arg Tyr Gly Gly Ile Ser Ser Ala Ser Asp Cys	
55		

285 290 295
 5 tct agt ctt cct tcc gca ctc caa gct ggt tgt aaa tgg aga ttc aac 1008
 Ser Ser Leu Pro Ser Ala Leu Gln Ala Gly Cys Lys Trp Arg Phe Asn
 300 305 310
 10 tgg ttc aag aac gct gat aac cca agc atg act tac aag gaa gtt acc 1056
 Trp Phe Lys Asn Ala Asp Asn Pro Ser Met Thr Tyr Lys Glu Val Thr
 315 320 325
 15 tgt ccc aag gaa atc acc gcc aag aca ggt tgt tca aga aaa taa 1101
 Cys Pro Lys Glu Ile Thr Ala Lys Thr Gly Cys Ser Arg Lys
 330 335 340
 20
 <210> 5
 25 <211> 360
 <212> PRT
 <213> Rhizopus oryzae CP96001
 30
 <220>
 <221> sig#peptide
 35 <222> (-23)... (-1)
 <220>
 40 <221> mat#peptide
 <222> (1)... (337)
 45
 <400> 5
 Met Lys Phe Leu Thr Ile Ala Ser Ser Ala Ile Leu Ala Leu Ala Val
 50 -20 -15 -10
 Gly Thr Glu Met Ala His Ala Ala Glu Cys Ser Lys Ala Tyr Tyr Gln
 -5 1 5
 55

Cys Gly Gly Lys Asn Trp Asp Gly Pro Thr Cys Cys Glu Ser Gly Ser
 10 15 20 25
 5 Thr Cys Val Asp Tyr Pro Asp Asn Pro Phe Tyr Ser Gln Cys Val Pro
 30 35 40
 10 Asn Glu Asn Leu Thr Ser Thr Asn Lys Ser Ser His Lys Thr Thr Thr
 45 50 55
 15 Thr Glu Ser Ala Lys Lys Thr Thr Thr Thr Lys Gly Ser Lys Lys Thr
 60 65 70
 Thr Thr Thr Glu Ala Ser Lys Lys Thr Thr Thr Thr Glu Ala Ser Lys
 75 80 85
 20 Lys Thr Thr Thr Thr Glu Ala Ser Lys Lys Thr Thr Thr Thr Thr Lys
 90 95 100 105
 25 Lys Ala Ser Thr Ser Thr Ser Ser Ser Ser Ser Ser Ala Ser Thr Asn
 110 115 120
 Tyr Ser Ala Val Ser Gly Gly Ala Ser Gly Asn Gly Glu Thr Thr Arg
 125 130 135
 30 Tyr Trp Asp Cys Cys Lys Pro Ser Cys Ser Trp Pro Gly Lys Ala Asp
 140 145 150
 35 Val Thr Ser Pro Val Gly Ser Cys Asn Lys Asp Gly Lys Thr Leu Ala
 155 160 165
 40 Asp Asn Asn Thr Gln Asn Gly Cys Val Gly Gly Ser Ser Tyr Thr Cys
 170 175 180 185
 Asn Asp Asn Gln Pro Trp Val Val Ser Asp Asp Leu Ala Tyr Gly Phe
 190 195 200
 45 Ala Ala Ala Ser Ile Ser Gly Gly Ser Glu Ala Thr Trp Cys Cys Ala
 205 210 215
 50 Cys Phe Glu Leu Thr Phe Thr Ser Thr Ala Val Lys Gly Lys Lys Met
 220 225 230
 55 Val Val Gln Val Thr Asn Thr Gly Ser Asp Leu Gly Ser Asn Thr Gly

235 240 245
 5 Ala His Phe Asp Leu Gln Met Pro Gly Gly Gly Val Gly Ile Tyr Asn
 250 255 260 265
 Gly Cys Ala Thr Gln Trp Gly Ala Pro Thr Asp Gly Trp Gly Ala Arg
 10 270 275 280
 Tyr Gly Gly Val Ser Ser Ala Ser Asp Cys Ser Asn Leu Pro Ser Ala
 285 290 295
 15 Leu Gln Ala Gly Cys Lys Trp Arg Phe Gly Trp Phe Lys Asn Ala Asp
 300 305 310
 20 Asn Pro Thr Met Thr Tyr Lys Gln Val Thr Cys Pro Lys Ala Ile Thr
 315 320 325
 Ala Lys Ser Gly Cys Ser Arg Lys
 25 330 335

 30 <210> 6
 <211> 1083
 <212> DNA
 <213> Rhizopus oryzae CP96001
 35
 <220>
 <221> sig#peptide
 <222> (1)... (69)
 40
 45 <220>
 <221> mat#peptide
 <222> (70)... (1083)
 50

 55 <400> 6
 atg aag ttc ctt acc att gcc tcc tcc gct atc ttg gca ctt gcc gtc 48

Met Lys Phe Leu Thr Ile Ala Ser Ser Ala Ile Leu Ala Leu Ala Val
 5 -20 -15 -10
 ggt act gaa atg gcc cat gct gct gaa tgt agc aag gct tac tac caa 96
 Gly Thr Glu Met Ala His Ala Ala Glu Cys Ser Lys Ala Tyr Tyr Gln
 10 -5 1 5
 tgt ggt ggt aag aac tgg gat gga cct acc tgc tgt gaa tct ggc tct 144
 Cys Gly Gly Lys Asn Trp Asp Gly Pro Thr Cys Cys Glu Ser Gly Ser
 15 10 15 20 25
 act tgc gtt gat tat cct gac aat cct ttc tac tcc caa tgt gtt ccc 192
 Thr Cys Val Asp Tyr Pro Asp Asn Pro Phe Tyr Ser Gln Cys Val Pro
 20 30 35 40
 aat gaa aac ctc acc tcc act aac aaa tct tct cac aaa acc acc act 240
 25 Asn Glu Asn Leu Thr Ser Thr Asn Lys Ser Ser His Lys Thr Thr Thr
 45 50 55
 act gag agt gcc aag aag act acc act act aaa ggt tcc aag aag acc 288
 30 Thr Glu Ser Ala Lys Lys Thr Thr Thr Thr Lys Gly Ser Lys Lys Thr
 60 65 70
 acc act act gaa gcc tct aag aag acc acc act act gaa gct tcc aag 336
 35 Thr Thr Thr Glu Ala Ser Lys Lys Thr Thr Thr Thr Glu Ala Ser Lys
 75 80 85
 aag acc acc act act gaa gcc tct aag aag acc acc act act act aag 384
 40 Lys Thr Thr Thr Thr Glu Ala Ser Lys Lys Thr Thr Thr Thr Thr Lys
 90 95 100 105
 45 aag gct tct acc tcc act tcc tct tcc tct tct tct gct tct aca aac 432
 Lys Ala Ser Thr Ser Thr Ser Ser Ser Ser Ser Ala Ser Thr Asn
 110 115 120
 50 tac tcc gct gtc tct ggt ggt gcc tcc ggt aat ggt gaa acc act cgc 480
 Tyr Ser Ala Val Ser Gly Gly Ala Ser Gly Asn Gly Glu Thr Thr Arg
 125 130 135
 55

5 tac tgg gat tgt tgt aag cct tct tgc agt tgg ccc ggt aag gct gat 528
 Tyr Trp Asp Cys Cys Lys Pro Ser Cys Ser Trp Pro Gly Lys Ala Asp
 140 145 150
 10 gtc acc tcc cct gtt ggc tcc tgt aac aag gat ggt aag act ctt gct 576
 Val Thr Ser Pro Val Gly Ser Cys Asn Lys Asp Gly Lys Thr Leu Ala
 155 160 165
 15 gat aac aac act caa aac ggc tgt gtt ggt ggt agc agc tac acc tgt 624
 Asp Asn Asn Thr Gln Asn Gly Cys Val Gly Gly Ser Ser Tyr Thr Cys
 170 175 180 185
 20 aat gac aat caa cct tgg gtt gtt agc gac gac ctt gcc tac ggt ttc 672
 Asn Asp Asn Gln Pro Trp Val Val Ser Asp Asp Leu Ala Tyr Gly Phe
 190 195 200
 25 gcc gct gct tcc att tct ggt ggt agc gaa gct act tgg tgt tgt gcc 720
 Ala Ala Ala Ser Ile Ser Gly Gly Ser Glu Ala Thr Trp Cys Cys Ala
 205 210 215
 30 tgt ttc gaa ctc aca ttc acc tct act gcc gtc aag ggt aag aag atg 768
 Cys Phe Glu Leu Thr Phe Thr Ser Thr Ala Val Lys Gly Lys Lys Met
 220 225 230
 35 gtt gtt caa gta acc aac act ggt tct gac ctt ggc tct aac act ggt 816
 Val Val Gln Val Thr Asn Thr Gly Ser Asp Leu Gly Ser Asn Thr Gly
 235 240 245
 40 gct cac ttt gac ttg caa atg ccc ggt ggt ggt gtt ggt atc tac aat 864
 Ala His Phe Asp Leu Gln Met Pro Gly Gly Gly Val Gly Ile Tyr Asn
 250 255 260 265
 45 ggt tgt gcc act caa tgg ggt gct ccc acc gat ggt tgg ggt gca aga 912
 Gly Cys Ala Thr Gln Trp Gly Ala Pro Thr Asp Gly Trp Gly Ala Arg
 270 275 280
 50 tac ggc ggt gtt tct tct gcc tct gac tgt tct aac ctt cct tct gcc 960
 Tyr Gly Gly Val Ser Ser Ala Ser Asp Cys Ser Asn Leu Pro Ser Ala
 55

	285	290	295	
5	ctt caa gct ggt tgt aag tgg aga ttc ggc tgg ttc aaa aac gct gat 1008 Leu Gln Ala Gly Cys Lys Trp Arg Phe Gly Trp Phe Lys Asn Ala Asp			
	300	305	310	
10	aac cca acc atg acc tac aaa caa gtt acc tgt ccc aag gct atc act 1056 Asn Pro Thr Met Thr Tyr Lys Gln Val Thr Cys Pro Lys Ala Ile Thr			
	315	320	325	
15	gcc aag tct ggc tgt tca aga aaa taa Ala Lys Ser Gly Cys Ser Arg Lys . 1083			
	330	335		
20				
	<210> 7			
25	<211> 338			
	<212> PRT			
	<213> Mucor circinelloides CP99001			
30				
	<220>			
	<221> sig#peptide			
35	<222> (-22)... (-1)			
	<220>			
40	<221> mat#peptide			
	<222> (1)... (316)			
45				
	<400> 7			
	Met Lys Phe Thr Val Ala Ile Thr Ser Ile Ala Val Ala Leu Ala Leu			
50	-20	-15	-10	
	Ser Ser Ser Ala Glu Ala Ala Ser Cys Ser Ser Val Tyr Gly Gln Cys			
	-5	1	5	10
55				

	Gly Gly Ile Gly Trp Ser Gly Pro Thr Cys Cys Glu Ser Gly Ser Thr
5	15 20 25
	Cys Val Ala Gln Glu Gly Asn Lys Tyr Tyr Ser Gln Cys Leu Pro Gly
	30 35 40
10	Ser His Ser Asn Asn Ala Gly Asn Ala Ser Ser Thr Lys Lys Thr Ser
	45 50 55
15	Thr Lys Thr Ser Thr Thr Thr Ala Lys Ala Thr Ala Thr Val Thr Thr
	60 65 70
	Lys Thr Val Thr Lys Thr Thr Thr Lys Thr Thr Thr Lys Thr Ser Thr
20	75 80 85 90
	Thr Ala Ala Ala Ser Thr Ser Thr Ser Ser Ser Ala Gly Tyr Lys Val
	95 100 105
25	Ile Ser Gly Gly Lys Ser Gly Ser Gly Ser Thr Thr Arg Tyr Trp Asp
	110 115 120
	Cys Cys Lys Ala Ser Cys Ser Trp Pro Gly Lys Ala Ser Val Thr Gly
30	125 130 135
	Pro Val Asp Thr Cys Ala Ser Asn Gly Ile Ser Leu Leu Asp Ala Asn
	140 145 150
35	Ala Gln Ser Gly Cys Asn Gly Gly Asn Gly Phe Met Cys Asn Asn Asn
	155 160 165 170
40	Gln Pro Trp Ala Val Asn Asp Glu Leu Ala Tyr Gly Phe Ala Ala Ala
	175 180 185
	Ser Ile Ala Gly Ser Asn Glu Ala Gly Trp Cys Cys Gly Cys Tyr Glu
45	190 195 200
	Leu Thr Phe Thr Ser Gly Ala Ala Ser Gly Lys Lys Met Val Val Gln
	205 210 215
50	Val Thr Asn Thr Gly Gly Asp Leu Gly Ser Asn His Phe Asp Leu Gln
	220 225 230
55	Met Pro Gly Gly Gly Val Gly Ile Phe Asn Gly Cys Ala Ala Gln Trp

235 240 245 250
 5 Gly Ala Pro Asn Asp Gly Trp Gly Ala Arg Tyr Gly Gly Val Ser Ser
 255 260 265
 Val Ser Asp Cys Ala Ser Leu Pro Ser Ala Leu Gln Ala Gly Cys Lys
 10 270 275 280
 Trp Arg Phe Asn Trp Phe Lys Asn Ser Asp Asn Pro Thr Met Thr Phe
 285 290 295
 15 Lys Glu Val Thr Cys Pro Ala Glu Leu Thr Thr Arg Ser Gly Cys Glu
 300 305 310
 20 Arg Lys
 315
 25 <210> 8
 <211> 1017
 <212> DNA
 30 <213> Mucor circinelloides CP99001
 <220>
 35 <221> sig#peptide
 <222> (1)... (66)
 40 <220>
 <221> mat#peptide
 45 <222> (67)... (1017)
 <400> 8
 50 atg aag ttc acc gtt gct att act tca atc gct gtt gca ctc gct ctc 48
 Met Lys Phe Thr Val Ala Ile Thr Ser Ile Ala Val Ala Leu Ala Leu
 -20 -15 -10
 55

EP 1 344 820 A1

5 agc tct tct gct gaa gct gct tct tgc agc tct gtc tat ggt caa tgt 96
 Ser Ser Ser Ala Glu Ala Ala Ser Cys Ser Ser Val Tyr Gly Gln Cys
 -5 1 5 10
 10 ggt ggc att gga tgg agt gga cct acc tgt tgt gaa agt ggc tct act 144
 Gly Gly Ile Gly Trp Ser Gly Pro Thr Cys Cys Glu Ser Gly Ser Thr
 15 15 20 25
 tgc gtt gct caa gaa ggc aac aaa tac tac tct caa tgt ctt ccc gga 192
 Cys Val Ala Gln Glu Gly Asn Lys Tyr Tyr Ser Gln Cys Leu Pro Gly
 30 35 40
 20 tcc cac agt aac aat gct ggt aac gct agc agc acc aag aag aca tct 240
 Ser His Ser Asn Asn Ala Gly Asn Ala Ser Ser Thr Lys Lys Thr Ser
 45 50 55
 25 acc aag aca tct act acc acc gcc aag gct act gct act gtc acc acc 288
 Thr Lys Thr Ser Thr Thr Thr Ala Lys Ala Thr Ala Thr Val Thr Thr
 60 65 70
 30 aag aca gta acc aag aca act acc aag aca act acc aag act agc act 336
 Lys Thr Val Thr Lys Thr Thr Thr Lys Thr Thr Thr Lys Thr Ser Thr
 75 80 85 90
 35 act gcc gct gct tct act tcc acc tct tct tct gct ggt tac aag gtc 384
 Thr Ala Ala Ala Ser Thr Ser Thr Ser Ser Ser Ala Gly Tyr Lys Val
 95 100 105
 40 atc tct ggc ggt aaa tct ggc agt ggt tcc aca act cgt tat tgg gat 432
 Ile Ser Gly Gly Lys Ser Gly Ser Gly Ser Thr Thr Arg Tyr Trp Asp
 110 115 120
 45 tgt tgt aaa gct tct tgc agc tgg cct gga aaa gct tct gtc act ggt 480
 Cys Cys Lys Ala Ser Cys Ser Trp Pro Gly Lys Ala Ser Val Thr Gly
 50 125 130 135
 cct gtt gac acc tgt gcc tcc aat ggt atc tct tta tta gat gcc aat 528
 Pro Val Asp Thr Cys Ala Ser Asn Gly Ile Ser Leu Leu Asp Ala Asn
 55

	140	145	150	
5	gct caa agt ggt tgt aac ggt ggt aat ggt ttc atg tgt aac aac aac			576
	Ala Gln Ser Gly Cys Asn Gly Gly Asn Gly Phe Met Cys Asn Asn Asn			
	155	160	165	170
10	caa cct tgg gct gtc aat gat gag ctc gct tac ggt ttc gct gct gcc			624
	Gln Pro Trp Ala Val Asn Asp Glu Leu Ala Tyr Gly Phe Ala Ala Ala			
	175	180	185	
15	tct att gct ggc tcc aac gaa gct gga tgg tgt tgt ggc tgt tat gaa			672
	Ser Ile Ala Gly Ser Asn Glu Ala Gly Trp Cys Cys Gly Cys Tyr Glu			
	190	195	200	
20	ttg acc ttc act tct ggc gct gct tct gga aag aag atg gtt gtt caa			720
	Leu Thr Phe Thr Ser Gly Ala Ala Ser Gly Lys Lys Met Val Val Gln			
	205	210	215	
25	glt acc aac acc ggt ggc gat tta ggc tct aac cac ttt gat ttg caa			768
	Val Thr Asn Thr Gly Gly Asp Leu Gly Ser Asn His Phe Asp Leu Gln			
30	220	225	230	
	atg ccc ggt ggt ggc gtt ggt atc ttc aat ggc tgt gct gct caa tgg			816
	Met Pro Gly Gly Gly Val Gly Ile Phe Asn Gly Cys Ala Ala Gln Trp			
35	235	240	245	250
	ggc gct ccc aat gat ggc tgg gga gct aga tat ggt ggt gtc agc tct			864
	Gly Ala Pro Asn Asp Gly Trp Gly Ala Arg Tyr Gly Gly Val Ser Ser			
40	255	260	265	
	gic tct gac tgt gcc tct ctt ccc tct gct ctt caa gct ggt tgt aaa			912
45	Val Ser Asp Cys Ala Ser Leu Pro Ser Ala Leu Gln Ala Gly Cys Lys			
	270	275	280	
	tgg aga ttc aac tgg ttc aag aac tct gat aac cct acc atg acc ttc			960
50	Trp Arg Phe Asn Trp Phe Lys Asn Ser Asp Asn Pro Thr Met Thr Phe			
	285	290	295	
55	aag gaa gtt acc tgt cct gct gaa tta act act cgc tca ggt tgc gaa			1008

Lys Glu Val Thr Cys Pro Ala Glu Leu Thr Thr Arg Ser Gly Cys Glu
300 305 310
aga aag taa
Arg Lys
315

<210> 9
<211> 387
<212> PRT
<213> Mucor circinelloides CP99001

<220>
<221> sig#peptide
<222> (-22)... (-1)

<220>
<221> mat#peptide
<222> (1)... (365)

<400> 9
Met Lys Phe Thr Val Ala Ile Thr Ser Ile Ala Val Ala Leu Ala Leu
-20 -15 -10
Ser Ser Ser Ala Glu Ala Ala Ser Cys Ser Ser Val Tyr Gly Gln Cys
-5 1 5 10
Gly Gly Ile Gly Trp Thr Gly Pro Thr Cys Cys Asp Ala Gly Ser Thr
15 20 25
Cys Lys Ala Gln Lys Asp Asn Lys Tyr Tyr Ser Gln Cys Ile Pro Lys
30 35 40
Pro Lys Gly Ser Ser Ser Ser Ser Ser Cys Ser Ser Val Tyr Ser Glu

	45	50	55
5	Cys Gly Gly Ile Gly Trp Ser Gly Pro Thr Cys Cys Glu Ser Gly Ser		
	60	65	70
	Thr Cys Val Ala Gln Glu Gly Asn Lys Tyr Tyr Ser Gln Cys Leu Pro		
10	75	80	85
	Gly Ser His Ser Asn Asn Ala Gly Asn Ala Ser Ser Thr Lys Lys Thr		
	95	100	105
15	Ser Thr Lys Thr Ser Thr Thr Thr Ala Lys Ala Thr Ala Thr Val Thr		
	110	115	120
	Thr Lys Thr Val Thr Lys Thr Thr Thr Lys Thr Thr Thr Lys Thr Ser		
20	125	130	135
	Thr Thr Ala Ala Ala Ser Thr Ser Thr Ser Ser Ala Gly Tyr Lys		
25	140	145	150
	Val Ile Ser Gly Gly Lys Ser Gly Ser Gly Ser Thr Thr Arg Tyr Trp		
	155	160	165
30	Asp Cys Cys Lys Ala Ser Cys Ser Trp Pro Gly Lys Ala Ser Val Thr		
	175	180	185
	Gly Pro Val Asp Thr Cys Ala Ser Asn Gly Ile Ser Leu Leu Asp Ala		
35	190	195	200
	Asn Ala Gln Ser Gly Cys Asn Gly Gly Asn Gly Phe Met Cys Asn Asn		
	205	210	215
40	Asn Gln Pro Trp Ala Val Asn Asp Glu Leu Ala Tyr Gly Phe Ala Ala		
	220	225	230
45	Ala Ser Ile Ala Gly Ser Asn Glu Ala Gly Trp Cys Cys Gly Cys Tyr		
	235	240	245
	Glu Leu Thr Phe Thr Ser Gly Ala Ala Ser Gly Lys Lys Met Val Val		
50	255	260	265
	Gln Val Thr Asn Thr Gly Gly Asp Leu Gly Ser Asn His Phe Asp Leu		
	270	275	280
55			

5 Gln Met Pro Gly Gly Gly Val Gly Ile Phe Asn Gly Cys Ala Ala Gln
 285 290 295
 10 Trp Gly Ala Pro Asn Asp Gly Trp Gly Ala Arg Tyr Gly Gly Val Ser
 300 305 310
 15 Ser Val Ser Asp Cys Ala Ser Leu Pro Ser Ala Leu Gln Ala Gly Cys
 315 320 325 330
 20 Lys Trp Arg Phe Asn Trp Phe Lys Asn Ser Asp Asn Pro Thr Met Thr
 335 340 345
 Phe Lys Glu Val Thr Cys Pro Ala Glu Leu Thr Thr Arg Ser Gly Cys
 350 355 360
 25 Glu Arg Lys
 365
 30 <210> 10
 <211> 1164
 <212> DNA
 <213> Mucor circinelloides CP99001
 35 <220>
 <221> sig#peptide
 <222> (1)... (66)
 40 <220>
 <221> mat#peptide
 <222> (67)... (1164)
 45 <400> 10
 atg aag ttc acc gtt gct att acf tca atc gct gtt gca ctc gct ctc 48
 Met Lys Phe Thr Val Ala Ile Thr Ser Ile Ala Val Ala Leu Ala Leu
 50 55

	-20	-15	-10	
5	agc tct tct gct gaa gct gct tct tgc agc tct gtc tat ggt caa tgt	96		
	Ser Ser Ser Ala Glu Ala Ala Ser Cys Ser Ser Val Tyr Gly Gln Cys			
	-5	1	5	10
10	ggc ggc att ggc tgg act ggt cct aca tgt tgt gat gct gga tgc acc	144		
	Gly Gly Ile Gly Trp Thr Gly Pro Thr Cys Cys Asp Ala Gly Ser Thr			
	15	20	25	
15	tgt aaa gct caa aag gat aac aaa tat tat tct caa tgt att ccc aaa	192		
	Cys Lys Ala Gln Lys Asp Asn Lys Tyr Tyr Ser Gln Cys Ile Pro Lys			
	30	35	40	
20	ccc aag ggt tcc tcc tca tca tca tca tgt agt tcc gtc tat agt caa	240		
	Pro Lys Gly Ser Ser Ser Ser Ser Ser Cys Ser Ser Val Tyr Ser Gln			
	45	50	55	
25	tgc ggt ggc att gga tgg agt gga cct acc tgt tgt gaa agt ggc tct	288		
	Cys Gly Gly Ile Gly Trp Ser Gly Pro Thr Cys Cys Glu Ser Gly Ser			
	60	65	70	
30	act tgc gtt gct caa gaa ggc aac aaa tac tac tct caa tgt ctt ccc	336		
	Thr Cys Val Ala Gln Glu Gly Asn Lys Tyr Tyr Ser Gln Cys Leu Pro			
	75	80	85	90
35	gga tcc cac agt aac aat gct ggt aac gct agc agc acc aag aag aca	384		
	Gly Ser His Ser Asn Asn Ala Gly Asn Ala Ser Ser Thr Lys Lys Thr			
	95	100	105	
40	tct acc aag aca tct act acc acc gcc aag gct act gct act gtc acc	432		
	Ser Thr Lys Thr Ser Thr Thr Thr Ala Lys Ala Thr Ala Thr Val Thr			
	110	115	120	
45	acc aag aca gla acc aag aca act acc aag aca act acc aag act agc	480		
	Thr Lys Thr Val Thr Lys Thr Thr Thr Lys Thr Thr Thr Lys Thr Ser			
	125	130	135	
50	act act gcc gct gct tct act tcc acc tct tct tct gct ggt tac aag	528		
55				

Thr Thr Ala Ala Ala Ser Thr Ser Thr Ser Ser Ser Ala Gly Tyr Lys
 140 145 150
 5 gtc atc tct ggc ggt aaa tct ggc agt ggt tcc aca act cgt tat tgg 576
 Val Ile Ser Gly Gly Lys Ser Gly Ser Gly Ser Thr Thr Arg Tyr Trp
 10 155 160 165 170
 gat tgt tgt aaa gct tct tgc agc tgg cct gga aaa gct tct gtc act 624
 Asp Cys Cys Lys Ala Ser Cys Ser Trp Pro Gly Lys Ala Ser Val Thr
 15 175 180 185
 ggt cct gtt gac acc tgt gcc tcc aat ggt atc tct tta tta gat gcc 672
 Gly Pro Val Asp Thr Cys Ala Ser Asn Gly Ile Ser Leu Leu Asp Ala
 20 190 195 200
 aat gct caa agt ggt tgt aac ggt ggt aat ggt ttc atg tgt aac aac 720
 Asn Ala Gln Ser Gly Cys Asn Gly Gly Asn Gly Phe Met Cys Asn Asn
 25 205 210 215
 aac caa cct tgg gct gtc aat gat gag ctc gct tac ggt ttc gct gct 768
 Asn Gln Pro Trp Ala Val Asn Asp Glu Leu Ala Tyr Gly Phe Ala Ala
 30 220 225 230
 gcc tct att gct ggc tcc aac gaa gct gga tgg tgt tgt ggc tgt tat 816
 Ala Ser Ile Ala Gly Ser Asn Glu Ala Gly Trp Cys Cys Gly Cys Tyr
 35 235 240 245 250
 gaa ttg acc ttc act tct ggc gct gct tct gga aag aag atg gtt gtt 864
 Glu Leu Thr Phe Thr Ser Gly Ala Ala Ser Gly Lys Lys Met Val Val
 40 255 260 265
 caa gtt acc aac acc ggt ggc gat tta ggc tct aac cac ttt gat ttg 912
 Gln Val Thr Asn Thr Gly Gly Asp Leu Gly Ser Asn His Phe Asp Leu
 45 270 275 280
 caa atg ccc ggt ggt ggc gtt ggt atc ttc aat ggc tgt gct gct caa 960
 Gln Met Pro Gly Gly Gly Val Gly Ile Phe Asn Gly Cys Ala Ala Gln
 50 285 290 295
 55

5 tgg ggc gct ccc aat gat ggc tgg gga gct aga tat ggt ggt gtc agc 1008
 Trp Gly Ala Pro Asn Asp Gly Trp Gly Ala Arg Tyr Gly Gly Val Ser
 300 305 310

10 tct gtc tct gac tgt gcc tct ctt ccc tct gct ctt caa gct ggt tgt 1056
 Ser Val Ser Asp Cys Ala Ser Leu Pro Ser Ala Leu Gln Ala Gly Cys
 315 320 325 330

15 aaa tgg aga ttc aac tgg ttc aag aac tct gat aac cct acc atg acc 1104
 Lys Trp Arg Phe Asn Trp Phe Lys Asn Ser Asp Asn Pro Thr Met Thr
 335 340 345

20 ttc aag gaa gtt acc tgt cct gct gaa tta act act cgc tca ggt tgc 1152
 Phe Lys Glu Val Thr Cys Pro Ala Glu Leu Thr Thr Arg Ser Gly Cys
 350 355 360

25 gaa aga aag taa 1164
 Glu Arg Lys
 365

30 <210> 11
 <211> 346

35 <212> PRT
 <213> Phycomyces nitens CP99002

40 <220>
 <221> sig#peptide

45 <222> (-19)... (-1)

50 <220>
 <221> mat#peptide
 <222> (1)... (327)

55

<400> 11

5	Met	Lys	Phe	Ser	Ile	Ile	Ala	Ser	Ala	Leu	Leu	Ala	Ala	Ser	Ser		
					-15					-10				-5			
10	Thr	Tyr	Ala	Ala	Glu	Cys	Ser	Gln	Gly	Tyr	Gly	Gln	Cys	Gly	Gly	Lys	
					1				5				10				
15	Met	Trp	Thr	Gly	Pro	Thr	Cys	Cys	Thr	Ser	Gly	Phe	Thr	Cys	Val	Gly	
					15				20				25				
20	Ala	Glu	Asn	Asn	Glu	Trp	Tyr	Ser	Gln	Cys	Ile	Pro	Asn	Asp	Gln	Val	
					30				35				40			45	
25	Gln	Gly	Asn	Pro	Lys	Thr	Thr	Thr	Thr	Thr	Thr	Thr	Lys	Ala	Ala	Thr	
						50					55				60		
30	Thr	Thr	Lys	Ala	Pro	Val	Thr	Thr	Thr	Lys	Ala	Thr	Thr	Thr	Thr	Thr	
						65				70				75			
35	Thr	Lys	Ala	Pro	Val	Thr	Thr	Thr	Lys	Ala	Thr	Thr	Thr	Thr	Thr	Thr	
						80				85				90			
40	Lys	Thr	Thr	Thr	Lys	Thr	Thr	Thr	Thr	Lys	Ala	Ala	Thr	Thr	Thr	Ser	
						95				100			105				
45	Ser	Ser	Asn	Thr	Gly	Tyr	Ser	Pro	Ile	Ser	Gly	Gly	Phe	Ser	Gly	Asn	
						110			115			120			125		
50	Gly	Arg	Thr	Thr	Arg	Tyr	Trp	Asp	Cys	Cys	Lys	Pro	Ser	Cys	Ala	Trp	
						130				135				140			
55	Asp	Gly	Lys	Ala	Ser	Val	Thr	Lys	Pro	Val	Leu	Thr	Cys	Ala	Lys	Asp	
						145				150				155			
60	Gly	Val	Ser	Arg	Leu	Gly	Ser	Asp	Val	Gln	Ser	Gly	Cys	Val	Gly	Gly	
						160				165			170				
65	Gln	Ala	Tyr	Met	Cys	Asn	Asp	Asn	Gln	Pro	Trp	Val	Val	Asn	Asp	Asp	
						175			180				185				
70	Leu	Ala	Tyr	Gly	Phe	Ala	Ala	Ala	Ser	Leu	Gly	Ser	Ala	Gly	Ala	Ser	
						190				195			200			205	

Ala Phe Cys Cys Gly Cys Tyr Glu Leu Thr Phe Thr Asn Thr Ala Val
210 215 220
Ala Gly Lys Lys Phe Val Val Gln Val Thr Asn Thr Gly Asp Asp Leu
225 230 235
Ser Thr Asn His Phe Asp Leu Gln Met Pro Gly Gly Gly Val Gly Tyr
240 245 250
Phe Asn Gly Cys Gln Ser Gln Trp Asn Thr Asn Thr Asp Gly Trp Gly
255 260 265
Ala Arg Tyr Gly Gly Ile Ser Ser Ile Ser Glu Cys Asp Lys Leu Pro
270 275 280 285
Thr Gln Leu Gln Ala Gly Cys Lys Trp Arg Phe Gly Trp Phe Lys Asn
290 295 300
Ala Asp Asn Pro Glu Val Thr Phe Lys Ala Val Thr Cys Pro Ala Glu
305 310 315
Ile Ile Ala Lys Thr Gly Cys Glu Arg Lys
320 325

<210> 12
<211> 1041
<212> DNA
<213> Phycomyces nitens CP99002

<220>
<221> sig#peptide
<222> (1)... (57)

<220>
<221> mat#peptide
<222> (58)... (1041)

5 <400> 12
 atg aag ttc tcc atc atc gct tcc gcc ctt ctc ctc gct gcc agc tcc 48
 Met Lys Phe Ser Ile Ile Ala Ser Ala Leu Leu Leu Ala Ala Ser Ser
 10 -15 -10 -5
 act tac gct gct gaa tgc agc caa ggc tat ggc cag tgt ggt ggc aag 96
 Thr Tyr Ala Ala Glu Cys Ser Gln Gly Tyr Gly Gln Cys Gly Gly Lys
 15 1 5 10
 atg tgg act ggt ccc acc tgc tgc acc tcc gcc ttc acc tgt gta ggt 144
 Met Trp Thr Gly Pro Thr Cys Cys Thr Ser Gly Phe Thr Cys Val Gly
 20 15 20 25
 gcc gaa aac aac gag tgg tac tct cag tgt atc ccc aac gat caa gtc 192
 Ala Glu Asn Asn Glu Trp Tyr Ser Gln Cys Ile Pro Asn Asp Gln Val
 25 30 35 40 45
 cag ggt aac ccc aag acc acc acc acc acc acc acc aag gct gcc act 240
 Gln Gly Asn Pro Lys Thr Thr Thr Thr Thr Thr Thr Thr Lys Ala Ala Thr
 30 50 55 60
 acc acc aag gct cct gtc acc acc acc aag gcc acc acc acc acc acc 288
 Thr Thr Lys Ala Pro Val Thr Thr Thr Thr Lys Ala Thr Thr Thr Thr Thr
 35 65 70 75
 acc aag gcc cct gtc acc acc acc aag gcc act act act acc acc acc 336
 Thr Lys Ala Pro Val Thr Thr Thr Thr Lys Ala Thr Thr Thr Thr Thr Thr
 40 80 85 90
 aag acc acc acc aag acc acc acc acc aag gct gcc acc acc acc tcc 384
 Lys Thr Thr Thr Lys Thr Thr Thr Thr Thr Lys Ala Ala Thr Thr Thr Ser
 45 95 100 105
 tct tcc aac act ggc tac agc ccc att tct ggt ggc ttc tct gga aac 432
 Ser Ser Asn Thr Gly Tyr Ser Pro Ile Ser Gly Gly Phe Ser Gly Asn
 50 110 115 120 125
 55

5 ggt cgc act acc cgc tac tgg gat tgc tgc aag ccc tct tgc gcc tgg 480
 Gly Arg Thr Thr Arg Tyr Trp Asp Cys Cys Lys Pro Ser Cys Ala Trp
 130 135 140
 10 gac gga aag gct tct gta act aag cct gta ctc acc tgt gcc aag gat 528
 Asp Gly Lys Ala Ser Val Thr Lys Pro Val Leu Thr Cys Ala Lys Asp
 145 150 155
 15 ggt gtc agc cgt ctc ggt tcc gat gtc cag agc ggt tgc gtc ggc ggc 576
 Gly Val Ser Arg Leu Gly Ser Asp Val Gln Ser Gly Cys Val Gly Gly
 160 165 170
 20 cag gcc tac atg tgc aat gac aac cag ccc tgg gtt gtc aat gac gac 624
 Gln Ala Tyr Met Cys Asn Asp Asn Gln Pro Trp Val Val Asn Asp Asp
 175 180 185
 25 ctt gcc tac ggt ttc gct gct gcc agt ctc ggt agc gcc ggt gcc tct 672
 Leu Ala Tyr Gly Phe Ala Ala Ala Ser Leu Gly Ser Ala Gly Ala Ser
 190 195 200 205
 30 gca ttc tgc tgc ggc tgt tac gag ctt acc ttc acc aac act gct gtc 720
 Ala Phe Cys Cys Gly Cys Tyr Glu Leu Thr Phe Thr Asn Thr Ala Val
 210 215 220
 35 gct ggc aag aag ttt gtc gtc cag gtc acc aac acc ggt gat gat ctc 768
 Ala Gly Lys Lys Phe Val Val Gln Val Thr Asn Thr Gly Asp Asp Leu
 225 230 235
 40 agc acc aac cac ttt gat ttg cag atg ccc ggc ggt ggt gtc ggc tac 816
 Ser Thr Asn His Phe Asp Leu Gln Met Pro Gly Gly Gly Val Gly Tyr
 240 245 250
 45 ttc aac ggc tgc cag tcc cag tgg aac acc aac acc gat ggc tgg ggt 864
 Phe Asn Gly Cys Gln Ser Gln Trp Asn Thr Asn Thr Asp Gly Trp Gly
 255 260 265
 50 gct cgc tat ggc ggt att agc tct att tca gag tgc gac aag ctt cct 912
 Ala Arg Tyr Gly Gly Ile Ser Ser Ile Ser Glu Cys Asp Lys Leu Pro
 55

270 275 280 285
 5 acc cag ttg cag gct ggt tgc aag tgg aga ttc gga tgg ttc aag aac 960
 Thr Gln Leu Gln Ala Gly Cys Lys Trp Arg Phe Gly Trp Phe Lys Asn
 290 295 300
 10 gct gac aac cca gag gtc acc ttc aag gct gtt act tgc cct gcc gag 1008
 Ala Asp Asn Pro Glu Val Thr Phe Lys Ala Val Thr Cys Pro Ala Glu
 305 310 315
 15 atc att gcc aag act ggt tgc gag cgc aag taa 1041
 Ile Ile Ala Lys Thr Gly Cys Glu Arg Lys
 320 325
 20

<210> 13

<211> 1043

<212> DNA

<213> Artificial Sequence

<220>

<223> Codon-optimized sequence corresponding to RCE I
 35 protein (SEQ. ID NO: 2)

<220>

<221> sig#peptide

<222> (16)... (84)

<220>

<221> mat#peptide

<222> (84)... (1043)

<400> 13

	ggatcctggg acaag atg aag ttc atc act atc gcc tcc tcc gcc ctc ctt	51		
5	Met Lys Phe Ile Thr Ile Ala Ser Ser Ala Leu Leu			
	-20	-15		
	gcc ctc gcc ctt ggc act gag atg gcc tcc gcc gct gag tgc tcc aag	99		
10	Ala Leu Ala Leu Gly Thr Glu Met Ala Ser Ala Ala Glu Cys Ser Lys			
	-10	-5	1	5
	ctc tac gga cag tgc ggc gga aag aac tgg aac ggc ccc acc tgc tgc	147		
15	Leu Tyr Gly Gln Cys Gly Gly Lys Asn Trp Asn Gly Pro Thr Cys Cys			
	10	15	20	
	gag agc ggc tgc acc tgc aag gtc tgc aat gac tac tac agc cag tgc	195		
20	Glu Ser Gly Ser Thr Cys Lys Val Ser Asn Asp Tyr Tyr Ser Gln Cys			
	25	30	35	
	ctg ccg agc ggc tcc tgc gga aac aag tgc agc gag tgc gcc cac aag	243		
25	Leu Pro Ser Gly Ser Ser Gly Asn Lys Ser Ser Glu Ser Ala His Lys			
	40	45	50	
	aag acc acg acc gct gcc cac aag aag acc acg acc gcc gct cac aag	291		
30	Lys Thr Thr Thr Ala Ala His Lys Lys Thr Thr Thr Ala Ala His Lys			
	55	60	65	
	aag act acg acc gct ccc gcc aag aag acc acg acc gtc gcc aag gct	339		
35	Lys Thr Thr Thr Ala Pro Ala Lys Lys Thr Thr Thr Val Ala Lys Ala			
	70	75	80	85
	tgc act ccg tcc aac tgc agc agc tgc tct tgc gga aag tac agc gct	387		
40	Ser Thr Pro Ser Asn Ser Ser Ser Ser Ser Ser Gly Lys Tyr Ser Ala			
	90	95	100	
	gtc agc ggt ggc gct agc ggc aac ggc gtc act acc cgc tac tgg gac	435		
45	Val Ser Gly Gly Ala Ser Gly Asn Gly Val Thr Thr Arg Tyr Trp Asp			
	105	110	115	
	tgc tgc aag gct tgc tgc tgc tgg ccc ggc aag gct aac gtc agc tgc	483		
50	Cys Cys Lys Ala Ser Cys Ser Trp Pro Gly Lys Ala Asn Val Ser Ser			

	120	125	130	
5	cct gtc aag tcc tgc aac aag gac ggc gtc acc gct ctt agc gac tcc			531
	Pro Val Lys Ser Cys Asn Lys Asp Gly Val Thr Ala Leu Ser Asp Ser			
	135	140	145	
10	aac gcc cag tcc ggc tgc aac ggc ggc aac tcc tac atg tgc aac gac			579
	Asn Ala Gln Ser Gly Cys Asn Gly Gly Asn Ser Tyr Met Cys Asn Asp			
	150	155	160	165
15	aac cag cca tgg gct gtc aac gac aac ctt gct tac ggt ttc gct gcc			627
	Asn Gln Pro Trp Ala Val Asn Asp Asn Leu Ala Tyr Gly Phe Ala Ala			
	170	175	180	
20	gct gcc att agc ggc ggt ggc gag agc cgc tgg tgc tgc tcc tgc ttc			675
	Ala Ala Ile Ser Gly Gly Gly Glu Ser Arg Trp Cys Cys Ser Cys Phe			
	185	190	195	
25	gag ctc acc ttc acc tcc acc agc gtt gct ggc aag aag atg gtc gtc			723
	Glu Leu Thr Phe Thr Ser Thr Ser Val Ala Gly Lys Lys Met Val Val			
	200	205	210	
30	cag gtc acc aac act ggc ggt gac ctt ggc agc tcg acc ggt gcc cac			771
	Gln Val Thr Asn Thr Gly Gly Asp Leu Gly Ser Ser Thr Gly Ala His			
	215	220	225	
35	ttc gat ctc cag atg ccc ggc ggc ggc gtc ggc atc ttc aac gga tgc			819
	Phe Asp Leu Gln Met Pro Gly Gly Gly Val Gly Ile Phe Asn Gly Cys			
	230	235	240	245
40	tcg tcc cag tgg ggc gct ccc aac gac ggc tgg ggc tcg cgc tac ggc			867
	Ser Ser Gln Trp Gly Ala Pro Asn Asp Gly Trp Gly Ser Arg Tyr Gly			
	250	255	260	
45	ggc atc agc tcc gcc agc gac tgc tcg tcc ctc ccc agc gcc ctc cag			915
	Gly Ile Ser Ser Ala Ser Asp Cys Ser Ser Leu Pro Ser Ala Leu Gln			
	265	270	275	
50	gcc ggc tgc aag tgg cgc ttc aac tgg ttc aag aac gcc gac aac ccg			963
55				

Ala Gly Cys Lys Trp Arg Phe Asn Trp Phe Lys Asn Ala Asp Asn Pro
 280 285 290
 5 tcc atg acc tac aag gag gtc acc tgc ccc aag gag atc acc gct aag 1011
 Ser Met Thr Tyr Lys Glu Val Thr Cys Pro Lys Glu Ile Thr Ala Lys
 10 295 300 305
 acc gga tgc tgc cgc aag taa acgcagg atcc 1043
 Thr Gly Cys Ser Arg Lys
 15 310 315
 <210> 14
 20 <211> 40
 <212> PRT
 25 <213> Rhizopus oryzae CP96001
 <400> 14
 30 Ala Glu Cys Ser Lys Leu Tyr Gly Gln Cys Gly Gly Lys Asn Trp Asn
 1 5 10 15
 Gly Pro Thr Cys Cys Glu Ser Gly Ser Thr Cys Lys Val Ser Asn Asp
 35 20 25 30
 Tyr Tyr Ser Gln Cys Leu Pro Ser
 35 40
 40
 <210> 15
 45 <211> 22
 <212> PRT
 <213> Mucor circinelloides CP99001
 50
 <400> 15
 Ala Ser Cys Ser Ser Val Tyr Gly Gln Cys Gly Gly Ile Gly Trp Ser
 55

1 5 10 15
 Gly Pro Thr Cys Cys Glu
 5 20

 10 <210> 16
 <211> 23
 <212> PRT
 15 <213> *Phycomyces nitens* CP99002

 <400> 16
 20 Ala Glu Cys Ser Gln Gly Tyr Gly Gln Cys Gly Gly Lys Met Trp Thr
 1 5 10 15
 Gly Pro Thr Cys Cys Thr Ser
 25 20

 30 <210> 17
 <211> 39
 <212> PRT
 35 <213> Artificial Sequence

 <220>
 40 <223> Consensus amino acid sequence found in cellulose binding
 domains of the family 45 endoglucanases

 45 <220>
 <221> Xaa
 50 <222> (1).. (6), (11).. (14), (16).. (18), (20).. (25), (27).. (31),
 (33).. (34), (36), (39)
 <223> Xaa is one of all amino acids
 55

5 <300>
 <301> Hoffren, A. -M. et al.
 <303> Protein Engineering
 10 <304> 8
 <306> 443-450
 <307> 1995
 15
 <400> 17
 Xaa Xaa Xaa Xaa Xaa Xaa Gln Cys Gly Gly Xaa Xaa Xaa Xaa Gly Xaa
 20 1 5 10 15
 Xaa Xaa Cys Xaa Xaa Xaa Xaa Xaa Xaa Cys Xaa Xaa Xaa Xaa Xaa Asn
 25 20 25 30
 Xaa Xaa Tyr Xaa Gln Cys Xaa
 35
 30
 <210> 18
 <211> 1257
 35 <212> DNA
 <213> Humicola insolens
 40
 <220>
 <221> intron
 <222> (453).. (509)
 45
 <400> 18
 50 aatgacgggg caacctcccg cccgggcca acctctgggt ttggttgac aggccgtctg 60
 tctcttgcgt cctcttacta cgcctgcctg gacctacgt ctcaactccg attcaagaig 120
 cgttctctcc ctctctcccg ctccgccgtt gtggccgcc tgcgggtgtt ggcccttgcc 180
 55

gctgatggca agtccaccog clactgggac tgcgtcaagc ctctgtcggg clgggccaag 240
 5 aaggctcccg tgaaccagcc tgccttctcc tgcaacgcca acttccagcg tctcactgac 300
 ttccagcgca agtccggctg cgagccgggc gggtcgcctt actcgtgcgc cgaccagacc 360
 ccatgggctg tgaacgacga ctctcgcttc gggtttgcig ccacctctat tgcgggcagc 420
 10 aatgaggcgg gctgggtgctg cgcttcttac gagtaagcct tggtcgcgtg tgtaacactg 480
 tgcaggcata gcactaacca cctcccaggc tcaccttacc atccggtcct gttgctggca 540
 agaagatggt cgtccagtc accagcactg gcggtgatct tggcagcaac cacttcgac 600
 15 tcaacatccc cggcggcggc gtcggcatct tgcacggtg cactccccag ttccggcgtc 660
 tgcccgccca gcgtacggc ggcatctcgt cccgcaacga gtgcgacgg tccccgacg 720
 cctcaagcc cggctgctac tggcgcttc acttggttcaa gaacgccgac aaccgagct 780
 20 tcagcttccg tcaggtccaa tgcacagccg agctcgtcgc tgcaccgga tgcggccgca 840
 acgacgacgg caacttccct gccgtccaga tcccctccag cagcaccagc tctccggtcg 900
 25 gccagcttac cagtaccagc accacctcca cctccaccac ctccagcccg cccgtccagc 960
 ctacgactcc cagcggctgc actgctgaga ggtgggctca gtccggcggc aatggctgga 1020
 gcggctgcac cacctgcgc gcgggcagca cctgcacgaa gattaatgac tggtaaccac 1080
 30 agtgcctgta aacgcagggc agcctgagaa ccttactggt tgcgcaacga aatgacactc 1140
 ccaatcacig tattagtct tttacataat ttctcatcc ctccagggat tgcacatat 1200
 atgcaatgat gaatactgaa cacaacactg gccgcttga cttggccgaag gaatgcc 1257

35 <210> 19

<211> 1720

40 <212> DNA

<213> *Trichoderma viride*

45 <220>

<221> intron

50 <222> (500).. (682)

55 <400> 19

gggtgigcat ttctccicaa catacigcct ttcaacaact ttgcctcct cctggcctg 60
 atatcccaat atcagttttt cccaagtag caagtaica gtaaatcgc tcatctatca 120
 5 ttaatcagtg cccatagtg ctcgtcttg atggcctccc gccatcacg atgaacagga 180
 ccatggctcc attgctgctt gcagcgtcga tactcttcgg gggcgtcgt gcacaacaga 240
 10 ctgtcgggg acagtgtaga ggtattggt ggagcggacc tacgattgt gtccttgat 300
 cagcttgct tactctcaat ccttattatg cgcaatgcat tccgggggcc actagtatca 360
 ccacctgac cggaccccc tgggtccaa ccaccaccac cagagccacc tcaacgacct 420
 15 catctccgc accgaccagc tctggagtc gatttgctgg cgttaacac gcgggctttg 480
 acttcggatg taccacagag tatgtcttca tgttgcatag tgttgctggc tgagtattct 540
 gggcggatga tttatagctg tgcggcgtgc aaaacaccgc cggctcgcca ctatcaaggc 600
 20 atagttgata ggcggcggtg tttcttcaa tccctgatt acactcicaa gaactagtg 660
 gctgaigat gtagtattac agtggcactt gcgttacac gaagtttat cctccgttga 720
 agaactcac tggggcaaac aactaccgg acggtatcgg ccagatgcag cacttcgtca 780
 25 acgatgaigg gatgactatt ttccgcctac ccgtcgatg gcagtacct gtaacaaca 840
 atctgggtg aactctcga tccaccagta tctcgaagta tgcagctc gttcaggggt 900
 gccgtctct cgtgtatc tgcacatcg acatccacaa ttatgctcga tggaacggtg 960
 30 gaalcatlgg ccaggaggg cctacaaatg cccagtttac cagcttttg tgcagttgg 1020
 catcgaagta cgcgtctcag tgcaggggt ggttcggaat aatgaatgag cccacgacg 1080
 35 tgaacatcaa cacttgggt gccacggtc aagaggctgt cactgcaac cgcaacgccg 1140
 gigtacgtc gcaatacat tctctgctg gaaatgatta tcaatctcg gcagctttta 1200
 ttccgatgg cagtgcagcc gccctgtctc aggtaacgaa cctgatgga tcaacaacga 1260
 40 atctaacttt cgatgtccac aagtacttag actcggacaa ctccggtact cagccgaat 1320
 gcactacaaa caacatcgac ggcgccttg ctccctcgc cacttggctt cgacagaaca 1380
 accgccagc tattctgacg gaaaccggcg gtggcaatgt tcagtcctgc atccaagatt 1440
 45 tgigccaaca gatccagtac ctcaaccaga actcagatgt ctatcttggc tatgttggct 1500
 ggggtgccg ttcatlgtat agcacttata ttctgacgga aagcctact ggaagcggt 1560
 50 actcgtggac ggacacatcc ctagtlagct cgtgtctgc caggaagtaa caccgaggtc 1620
 gattgcagga gccctgtcaa tagcatctt atcttctgt acataattct tactctctga 1680
 agccgcttgt tctgggtatg tctcttgaca ggttcttga 1720

5 <210> 20
 <211> 20
 <212> DNA
 10 <213> Artificial Sequence

 <220>
 15 <223> Description of Artificial Sequence:Primer

 <400> 20
 20 caccacgcgc tactgggact 20

 25 <210> 21
 <211> 19
 <212> DNA
 30 <213> Artificial Sequence

 <220>
 35 <223> Description of Artificial Sequence:Primer

 40 <400> 21
 ggatccctgcg ttacttgc 19

 45 <210> 22
 <211> 18
 <212> DNA
 50 <213> Artificial Sequence

<220>

5 <223> Description of Artificial Sequence:Primer

<400> 22

10 ggatcctggg acaagatg 18

<210> 23

15 <211> 17

<212> DNA

20 <213> Artificial Sequence

<220>

25 <223> Description of Artificial Sequence:Primer

<400> 23

30 gcacgacggc ttgcagc 17

<210> 24

35 <211> 228

<212> PRT

40 <213> Artificial Sequence

<220>

45 <223> Description of Artificial Sequence:Recombinant
protein

50 <400> 24

Met Gln Leu Pro Leu Thr Thr Leu Leu Thr Leu Leu Pro Ala Leu Ala

55 1 5 10 15

5	Ala Ala Gln Ser Gly Ser Gly Arg Thr Thr Arg Tyr Trp Asp Cys Cys	20	25	30
10	Lys Pro Ser Cys Ser Trp Pro Gly Lys Ala Asn Val Ser Ser Pro Val	35	40	45
15	Lys Ser Cys Asn Lys Asp Gly Val Thr Ala Leu Ser Asp Ser Asn Ala	50	55	60
20	Gln Ser Gly Cys Asn Gly Gly Asn Ser Tyr Met Cys Asn Asp Asn Gln	65	70	75
25	Pro Trp Ala Val Asn Asp Asn Leu Ala Tyr Gly Phe Ala Ala Ala Ala	85	90	95
30	Ile Ser Gly Gly Gly Glu Ser Arg Trp Cys Cys Ser Cys Phe Glu Leu	100	105	110
35	Thr Phe Thr Ser Thr Ser Val Ala Gly Lys Lys Met Val Val Gln Val	115	120	125
40	Thr Asn Thr Gly Gly Asp Leu Gly Ser Ser Thr Gly Ala His Phe Asp	130	135	140
45	Leu Gln Met Pro Gly Gly Gly Val Gly Ile Phe Asn Gly Cys Ser Ser	145	150	155
50	Gln Trp Gly Ala Pro Asn Asp Gly Trp Gly Ser Arg Tyr Gly Gly Ile	160		

	165	170	175
5	Ser Ser Ala Ser Asp Cys Ser Ser Leu Pro Ser Ala Leu Gln Ala Gly		
	180	185	190
10	Cys Lys Trp Arg Phe Asn Trp Phe Lys Asn Ala Asp Asn Pro Ser Met		
	195	200	205
15	Thr Tyr Lys Glu Val Thr Cys Pro Lys Glu Ile Thr Ala Lys Thr Gly		
20	210	215	220
	Cys Ser Arg Lys		
25	225		
30	<210> 25		
	<211> 20		
	<212> DNA		
35	<213> Artificial Sequence		
40	<220>		
	<223> Description of Artificial Sequence:Primer		
45	<400> 25		
	gcggatcctg ggacaagatg		20
50	<210> 26		
	<211> 24		
55	<212> DNA		

<213> Artificial Sequence

5

<220>

<223> Description of Artificial Sequence:Primer

10

<400> 26

gcctgcagag cggcggagc catc

24

15

<210> 27

<211> 25

20

<212> DNA

<213> Artificial Sequence

25

<220>

<223> Description of Artificial Sequence:Primer

30

<400> 27

gcctgcaggg aaagtacagc gctgt

25

35

<210> 28

<211> 21

40

<212> DNA

<213> Artificial Sequence

45

<220>

<223> Description of Artificial Sequence:Primer

50

<400> 28

gcggatccctg cgattacttg c

21

55

<210> 29

<211> 245

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Recombinant protein

<400> 29

Met Lys Phe Ile Thr Ile Ala Ser Ser Ala Leu Leu Ala Leu Ala Leu

1

5

10

15

Gly Thr Glu Met Ala Ser Ala Ala Leu Gln Gly Lys Tyr Ser Ala Val

20

25

30

Ser Gly Gly Ala Ser Gly Asn Gly Val Thr Thr Arg Tyr Trp Asp Cys

35

40

45

Cys Lys Pro Ser Cys Ser Trp Pro Gly Lys Ala Asn Val Ser Ser Pro

50

55

60

Val Lys Ser Cys Asn Lys Asp Gly Val Thr Ala Leu Ser Asp Ser Asn

65

70

75

80

Ala Gln Ser Gly Cys Asn Gly Gly Asn Ser Tyr Met Cys Asn Asp Asn

85

90

95

5	Gln Pro Trp Ala Val Asn Asp Asn Leu Ala Tyr Gly Phe Ala Ala Ala	100	105	110	
10	Ala Ile Ser Gly Gly Gly Glu Ser Arg Trp Cys Cys Ser Cys Phe Glu	115	120	125	
15	Leu Thr Phe Thr Ser Thr Ser Val Ala Gly Lys Lys Met Val Val Gln	130	135	140	
20	Val Thr Asn Thr Gly Gly Asp Leu Gly Ser Ser Thr Gly Ala His Phe	145	150	155	160
25	Asp Leu Gln Met Pro Gly Gly Gly Val Gly Ile Phe Asn Gly Cys Ser	165	170	175	
30	Ser Gln Trp Gly Ala Pro Asn Asp Gly Trp Gly Ser Arg Tyr Gly Gly	180	185	190	
35	Ile Ser Ser Ala Ser Asp Cys Ser Ser Leu Pro Ser Ala Leu Gln Ala	195	200	205	
40					
45	Gly Cys Lys Trp Arg Phe Asn Trp Phe Lys Asn Ala Asp Asn Pro Ser	210	215	220	
50	Met Thr Tyr Lys Glu Val Thr Cys Pro Lys Glu Ile Thr Ala Lys Thr	225	230	235	240
55	Gly Cys Ser Arg Lys	245			

5
10
15
20
25
30
35
40
45
50
55

<210> 30
<211> 7
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:N-terminus
sequence of recombinant protein

<400> 30
Gln Ser Gly Ser Gly Arg Thr
1 5

<210> 31
<211> 7
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:N-terminus
sequence of recombinant protein

<400> 31
Lys Tyr Ser Ala Val Ser Gly
1 5

<210> 32

<211> 7

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:N-terminus
sequence of recombinant protein

<400> 32

Ala Val Ser Gly Gly Ala Ser

1 5

<210> 33

<211> 7

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:N-terminus
sequence of recombinant protein

<400> 33

Ser Ala Val Ser Gly Gly Ala

1 5

5 <210> 34
 <211> 7
 <212> PRT
 <213> Artificial Sequence
 10
 <220>
 <223> Description of Artificial Sequence:N-terminus
 15 sequence of recombinant protein

 <400> 34
 Gly Gly Ala Ser Gly Asn Gly
 1 5
 25

 <210> 35
 30 <211> 7
 <212> PRT
 <213> Artificial Sequence
 35
 <220>
 <223> Description of Artificial Sequence:N-terminus
 40 sequence of recombinant protein

 <400> 35
 Ala Glu Cys Ser Lys Leu Tyr
 1 5
 50

 <210> 36
 55

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Primer

<400> 36

taytgggayt gytgyaarcc

20

<210> 37

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Primer

 $\langle 220 \rangle$

<221> modified#base

<222> (3)

<223> i

<220>

<221> modified#base

<222> (9)

<223> i

<400> 37

tcngcrtina rraaccaric

20

5

<210> 38

10

<211> 223

<212> PRT

<213> Humicola insolens

15

<400> 38

20

Met Gln Leu Pro Leu Thr Thr Leu Leu Thr Leu Leu Pro Ala Leu Ala

1 5 10 15

25

Ala Ala Gln Ser Gly Ser Gly Arg Thr Thr Arg Tyr Trp Asp Cys Cys

20 25 30

30

Lys Pro Ser Cys Ala Trp Pro Gly Lys Gly Pro Ala Pro Val Arg Thr

35 40 45

35

Cys Asp Arg Trp Asp Asn Pro Leu Phe Asp Gly Gly Asn Thr Arg Ser

50 55 60

40

Gly Cys Asp Ala Gly Gly Gly Ala Tyr Met Cys Ser Asp Gln Ser Pro

65 70 75 80

45

Trp Ala Val Ser Asp Asp Leu Ala Tyr Gly Trp Ala Ala Val Asn Ile

85 90 95

50

Ala Gly Ser Asn Glu Arg Gln Trp Cys Cys Ala Cys Tyr Glu Leu Thr

55

100 105 110

5 Phe Thr Ser Gly Pro Val Ala Gly Lys Arg Met Ile Val Gln Ala Ser
115 120 125

10 Asn Thr Gly Gly Asp Leu Gly Asn Asn His Phe Asp Ile Ala Met Pro
130 135 140

15 Gly Gly Gly Val Gly Ile Phe Asn Ala Cys Thr Asp Gln Tyr Gly Ala
145 150 155 160

20 Pro Pro Asn Gly Trp Gly Gln Arg Tyr Gly Gly Ile Ser Gln Arg His
165 170 175

25 Glu Cys Asp Ala Phe Pro Glu Lys Leu Lys Pro Gly Cys Tyr Trp Arg
180 185 190

30 Phe Asp Trp Phe Leu Asn Ala Asp Asn Pro Ser Val Asn Trp Arg Gln
195 200 205

35 Val Ser Cys Pro Ala Glu Ile Val Ala Lys Ser Gly Cys Ser Arg
40 210 215 220

45 <210> 39

<211> 672

<212> DNA

50

<213> Humicola insolens

55 <400> 39

atgcagctcc cccctgaccac gctccctcacc ctccctccccg cccctcgcgge ggcccagctcc 60
 ggcagcggcc gcaccacgcg ctactgggac tgcctgaagc cgtcgtgcgc gtggcccggc 120
 aagggcccgg cggccgtgcg gacgtgcgac cggctgggaca acccgctggt cgacggcgge 180
 aacacgcgca gcgggagcga cgcggcgggc ggcgcctaca tglgctcgga ccagagcccc 240
 tgggcgggca gcgacgacct ggcgtacggc tggcgggcgg tcaacattgc cggctccaac 300
 gagaggcagt ggtgctgcgc ctgctacgag ctgaccttca ccagcgggcc ggtggcgggc 360
 aagaggatga ttgtgcaggc gagcaacacg ggaggcgatt tggggaacaa ccactttgat 420
 attgctatgc cggcggtggc cgtcggctac ttcaacgcct gcaccgacca gtacggcgcg 480
 ccccccaacg gctggggcca gcgctacggc ggcatcagcc aacgccacga gtgcgacgcc 540
 ttccccgaga agctcaagcc cggctgctac tggcgcttgg aciggttccct caacgccgac 600
 aaccgagcgc tcaactggcg gcaggtcagc tggccggcgg agattgtggc caagagcggc 660
 tgctcgcggt aa 672

<210> 40

<211> 38

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Primer

<400> 40

ggggatccctg ggacaagatg cagctcccc tgaccacg

38

<210> 41

<211> 38

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Primer

5

<400> 41

10

ggggatccg catttaacgc gagcagccgc tcttgcc

38

<210> 42

15

<211> 21

<212> DNA

<213> Artificial Sequence

20

<220>

<223> Description of Artificial Sequence:Primer

25

<400> 42

30

gactgctgca agccgtcgtg c

21

<210> 43

35

<211> 20

<212> DNA

<213> Artificial Sequence

40

<220>

<223> Description of Artificial Sequence:Primer

45

<400> 43

50

gllgcacatg taggagttgc

20

55

Claims

1. A protein that is a *Zygomycetes*-derived endoglucanase lacking the cellulose-binding domain and exhibits endoglucanase activity.
2. A protein that is a *Zygomycetes*-derived endoglucanase belonging to family 45 lacking the cellulose-binding domain and exhibits endoglucanase activity.
3. The protein according to claim 1 or 2, wherein the *Zygomycetes* are microorganisms selected from the group consisting of those belonging to *Rhizopus*, *Mucor*, and *Phycomyces*.
4. The protein according to claim 3, wherein the *Zygomycetes* are microorganisms belonging to *Rhizopus*.
5. A protein comprising an amino acid sequence as shown in SEQ ID NO: 1, 3, 5, 7, 9, or 11, wherein the cellulose-binding domain has been deleted therefrom, and exhibiting endoglucanase activity, a modified protein thereof exhibiting endoglucanase activity, or a homologue of the protein or the modified protein exhibiting endoglucanase activity.
6. A protein comprising an amino acid sequence as shown in SEQ ID NO: 1, 3, 5, 7, 9, or 11, wherein the cellulose-binding domain has been deleted therefrom, and exhibiting endoglucanase activity.
7. A gene encoding the protein, modified protein thereof, or homologue of the protein or the modified protein according to any one of claims 1 to 6.
8. An expression vector comprising the gene according to claim 7.
9. A host cell transformed with the expression vector according to claim 8.
10. The host cell according to claim 9, which is a filamentous fungus.
11. The host cell according to claim 10, which is a microorganism belonging to *Humicola*.
12. A method for producing a protein comprising steps of culturing the host cell according to any one of claims 9 to 11 and collecting from the host cell obtained by the step of culturing or its culture product the protein, modified protein thereof, or homologue of the protein or the modified protein according to any one of claims 1 to 6.
13. A protein produced by the method according to claim 12.
14. A cellulase preparation comprising the protein, modified protein thereof, or homologue of the protein or the modified protein according to any one of claims 1 to 6 and 13.
15. A method for treating cellulose-containing fabrics comprising a step of bringing cellulose-containing fabrics into contact with the protein, modified protein thereof, or homologue of the protein or the modified protein according to any one of claims 1 to 6 and 13 or the cellulase preparation according to claim 14.
16. A method for reducing the rate at which cellulose-containing fabrics become fuzzy or reducing fuzzing in cellulose-containing fabrics comprising a step of bringing cellulose-containing fabrics into contact with the protein, modified protein thereof, or homologue of the protein or the modified protein according to any one of claims 1 to 6 and 13 or the cellulase preparation according to claim 14.
17. A method of weight loss treatment of cellulose-containing fabrics to improve the feel and appearance thereof comprising a step of bringing cellulose-containing fabrics into contact with the protein, modified protein thereof, or homologue of the protein or the modified protein according to any one of claims 1 to 6 and 13 or the cellulase preparation according to claim 14.
18. A method of color clarification of colored cellulose-containing fabrics comprising a step of treating colored cellulose-containing fabrics with the protein, modified protein thereof, or homologue of the protein or the modified protein according to any one of claims 1 to 6 and 13 or the cellulase preparation according to claim 14.

19. A method of providing colored cellulose-containing fabrics with partial color change comprising a step of treating colored cellulose-containing fabrics with the protein, modified protein thereof, or homologue of the protein or the modified protein according to any one of claims 1 to 6 and 13 or the cellulase preparation according to claim 14.
- 5 20. A method for reducing the rate at which cellulose-containing fabrics become stiff or reducing stiffness in cellulose-containing fabrics comprising a step of treating cellulose-containing fabrics with the protein, modified protein thereof, or homologue of the protein or the modified protein according to any one of claims 1 to 6 and 13 or the cellulase preparation according to claim 14.
- 10 21. The method according to any one of claims 15 to 20, wherein fabrics are treated through the soaking, washing, or rinsing thereof.
- 15 22. An additive to detergent comprising the protein, modified protein thereof, or homologue of the protein or the modified protein according to any one of claims 1 to 6 and 13 or the cellulase preparation according to claim 14 in a non-dusting granular form or a stabilized liquid form.
- 20 23. A detergent composition comprising the protein, modified protein thereof, or homologue of the protein or the modified protein according to any one of claims 1 to 6 and 13 or the cellulase preparation according to claim 14.
- 25 24. A method of deinking waste paper using a deinking agent wherein the protein, modified protein thereof, or homologue of the protein or the modified protein according to any one of claims 1 to 6 and 13 or the cellulase preparation according to claim 14 is used in a step of deinking waste paper.
- 25 25. A method for improving drainage of paper pulp comprising a step of treating paper pulp with the protein, modified protein thereof, or homologue of the protein or the modified protein according to any one of claims 1 to 6 and 13 or the cellulase preparation according to claim 14.
- 30 26. A method for improving digestibility of animal feeds comprising a step of treating animal feeds with the protein, modified protein thereof, or homologue of the protein or the modified protein according to any one of claims 1 to 6 and 13 or the cellulase preparation according to claim 14.

35

40

45

50

55

INTERNATIONAL SEARCH REPORT

International application No.

PCT/JP01/10188

A. CLASSIFICATION OF SUBJECT MATTER

Int.Cl.⁷ C12N 15/56, 1/15, 9/42, C11D 3/386, D06M 16/00 // (C12N 15/56, C12R 1:845), (C12N 15/56, C12R 1:645), (C12N 15/56, C12R 1:785), (C12N 1/15, C12R 1:645), (C12N 9/42, C12R 1:645), D06M101:06

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

Int.Cl.⁷ C12N 15/00-15/90, 9/42

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

MEDLINE (STN), WPI (DIALOG), BIOSIS (DIALOG), GenBank/EMBL/DBJ/GeneSeq, SwissProt/PIR/GeneSeq

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	AZEVEDO, H. et al., "Effects of agitation level on the adsorption, desorption, and activities on cotton fabrics of full length and core domains of EGV (<i>Humicola insolens</i>) and CenA (<i>Cellulomonas fimi</i>).", Enzyme Microb. Technol., August, 2000, Vol.27, No.3-5, pages 325 to 329	1-26
Y	WO 00/24879 A1 (Meiji Seika Kaisha, Ltd.), 04 May, 2000 (04.05.2000), & AU 9962300 A & EP 1123974 A1	1-26
P,A	WO 01/90375 A1 (Meiji Seika Kaisha, Ltd.), 29 November, 2001 (29.11.2001), (Family: none)	1-26
A	WO 94/21801 A2 (Genencor International, Inc.), 29 September, 1994 (29.09.1994), & FI 9504330 A & US 5475101 A & EP 689598 A1 & US 5753484 A	1-26

☒ Further documents are listed in the continuation of Box C.
 ☐ See patent family annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"I" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search
07 February, 2002 (07.02.02)Date of mailing of the international search report
26 February, 2002 (26.02.02)Name and mailing address of the ISA/
Japanese Patent Office

Authorized officer

Facsimile No.

Telephone No.

Form PCT/ISA/210 (second sheet) (July 1992)

INTERNATIONAL SEARCH REPORT

International application No.

PCT/JP01/10188

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	TAKASHIMA, S. et al., "Comparison of gene structures and enzymatic properties between two endoglucanases from <i>Humicola grisea</i> ", J. Biotechnol., (1999), Vol.67, No.2-3, pages 85 to 97	1-26

Form PCT/ISA/210 (continuation of second sheet) (July 1992)